

A gene encoding a protein, nestin, whose expression distinguishes neural multipotential stem cells and brain tumor cells from the more differentiated neural cell types (e.g., neuronal, glial and muscle cells). Methods of detecting the expression of nestin as a means of diagnosing and treating brain tumours are also described.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

-1-

NESTIN EXPRESSION AS AN INDICATOR OF  
NEUROEPITHELIAL TUMORS

Description

Background

05       Brain tumors are a leading cause of cancer deaths in people under the age of 35. An important concept in our understanding of brain tumors is that the neoplastic cells may arise from undifferentiated neuroectodermal cells (Rubinstein, L.J. J. Neurosurg 62: 795-805 (1985)).

10       At the gastrula stage of central nervous system (CNS) development, neuroectoderm, forming the neural plate is derived from ectoderm which has been induced by underlying mesoderm. (Spemann, H. Yale University Press 1938)). The neural plate then folds, in the process  
15       called neurulation, to form the neural tube. The cells of the neural tube are the precursors for the major differentiated cell types in the central nervous system: astrocytes, oligodendrocytes, and the various types of neuron (Sauer, F.C. J. Comp. Neurol. 62: 377-405  
20       (1935)). The cells of the neural tube may also be the precursors to neoplastic cells which form brain tumors.

      Recent success in "tagging" individual CNS pre-cursors and their daughter cells, either through infection by a marker-bearing retrovirus or by injection of  
25       stable, specific dyes and enzymes, has shown that neurons and non-neuronal cells (i.e., glia cells) are often derived from a common precursor. In retina a common precursor for neurons and the Mueller glial cell exists

-2-

very late in development (Holt, C.E., T.W. Bertsch et al., Neuron 1: 15-26 (1988); Turner, D.L. and C.L. Cepko, Nature 238: 131-136 (1987); Wetts, R. and S.E. Fraser, Science 239: 1142-1145 (1988); a common precursor for  
05 different neuronal and glial cell types has also been demonstrated in cortex and tectum (Luskin, M.B., A.L. Pearlman et al., Neuron 1: 635-647 (1988); Price, J. and L. Thurlow, Development 104: 473-482 (1988); Walsh, C. and C. Cepko, Science 241: 1342-1345 (1988)).

10 Tissue culture systems have provided additional information about the differentiation potential of CNS stem cells. Single-cell microculture of embryonic day 14 rat forebrain has produced clones containing both neurons and glia (Temple, S. Nature 340: 471-473 (1989)).

15 Analysis of a later precursor in the glial differentiation pathway has established specific factors influencing differentiation to oligodendrocytes or type II astrocytes (Raff, M. Science 243: 1450-1455 (1989)). Cell  
20 lines established by immortalization of CNS stem cells also reflect features of the stem cell and can differentiate along both the neuronal and glial pathways (Cepko, C.L. Ann. Rev. Neurosci. 12: 47-65 (1989); Frederiksen, K. et al., supra (1988)).

Antibodies have also proven useful for analyzing  
25 stem cells. For example, antibodies A2B5 (Raff, M.C. supra 1989) D1.1 (Levine, J.M., L. Beasley et al., J. Neurosci 4: 820-831 (1984), Rat 401 (Hockfield, S. and R. McKay, J. Neurosci 5: 3310-3328 (1985), and antisera  
30 against the intermediate filament vimentin (Bignami, A., T. Raju et al Dev. Biol 91: 286-295 (1982); Federoff, S.

-3-

Molecular Bases of Neural Development (1985); Tapscott, S.J., G.S. Bennett et al., Dev. Biol. 86: 40-54 (1981) bind to antigens enriched in the proliferative zone of the neural tube. However, A2B5 and D1.1 recognize glycolipid epitopes and are less useful for analysis of gene expression. Vimentin, which appears transiently in brain development, is an ambiguous marker largely because of its promiscuous expression in many cultured cells as well as in a variety of developing and differentiated tissues (Traub, N. Intermediate filaments: a review. Berlin, SpringerVerlag (1985)).

The correlation between Rat 401 and proliferating cells in the developing nervous system has been examined in detail. Rat 401 was found to recognize a transient population of embryonic columnar epithelial cells and radial glial cells in many regions of the CNS (Hockfield, S. and McKay J. J. Neurosci. 5: 3310-3328 (1985)). The number of Rat 401 positive cells, their proliferative rate, and the developmental kinetics of Rat 401 expression relative to neuronal differentiation revealed that the immediate precursors to neurons are also Rat 401 positive (Frederiksen and McKay, J. J. Neurosci 8: 1144-1151 (1988)). Neither neurons nor glia in the adult brain express the epitope recognized by the Rat 401 antibody.

Although "tagging", tissue culture systems and antibodies have proven useful in analyzing neural precursor cells, there is still an incomplete understanding of the lineage of tumor development. For example, medulloblastoma, a common brain tumor of children, is associated predominantly with the cerebellum and brain

-4-

stem. Medulloblastoma tumors contain cell types with differentiated characteristics of neurons, glia and muscle (Rubinstein, supra 1985; Coakham H.B., et al., J. Clin. Pathol. 38: 165-173 (1985); Velasco, M.E. et al., 05 Surg. Neurol. 23: 177-182 (1985); Hayashi, K. et al., Acta Pathologica Japonica 37: 85-96 (1987); Cras, P. et al., Acta. Neuropathol 75: 377-384 (1988)).

Because of the presence of multiple differentiated cell types found in these tumors, Bailey and Cushing 10 (Bailey, P. and H. Cushing, Arch. Neurol. Psychiatry 14: 192-224 (1925) proposed that rather than being multiple coincident tumors, they were derived from a multi-potential stem cell, which they called a "medulloblast". According to this model, a medulloblastoma tumor would 15 result from uncontrolled proliferation and differentiation of the medulloblast. However, the hypothetical medulloblast has not been identified.

Due to the incomplete understanding of brain tumor development, there is currently no adequate assay for 20 detecting brain tumors at an early stage in their development. In addition, there is no specific noninvasive method of treating brain tumors. Methods of detecting and treating brain tumors would be very useful.

#### Summary of the Invention

25 The present invention relates to a gene encoding a protein whose expression distinguishes neural multi-potential stem cells and brain tumor cells from the more differentiated neural cell types (e.g. neuronal, glial and muscle cells) of the mammalian brain.

-5-

The present invention relates to the nestin gene, particularly a nestin gene of mammalian origin, and the encoded nestin protein. As described herein, two nestin genes of mammalian origin, the rat nestin gene and the human nestin gene, have been isolated and sequenced. The coding sequence of the rat nestin gene transcript is 5415 bp, which corresponds to an expected molecular weight of approximately 200 kD for the nestin protein. The relative molecular weight of the nestin protein as determined by SDS polyacrylamide gels is approximately 240 kD. The predicted amino acid sequence of the rat nestin gene product shows that nestin defines a distinct sixth class of intermediate filament protein.

The present invention also relates to the human nestin gene and the encoded human nestin protein. The coding sequence of the human nestin gene transcript is 4854 bp. The predicted amino acid sequence is 1618 amino acids in length, and shows significant sequence homology with the rat nestin sequence.

The present invention further relates to methods of detecting the expression of nestin as a means of diagnosing a predisposition to the development of a brain tumor or the presence of a brain tumor in an adult individual. In one embodiment, DNA present in a sample from the brain of an adult individual is hybridized to a DNA probe which is complementary to all or a portion of the nestin gene. As used herein, the term the nestin gene includes the human nestin gene whose sequence is represented herein, the rat nestin gene whose sequence is also represented herein and equivalent genes from other

-6-

species, such as those which are substantially homologous to the human or the rat gene sequence. Detection of hybridization is an indication of a predisposition to the development of a brain tumor or the presence of a brain  
05 tumor. In another embodiment, cerebral spinal fluid or a serum sample from the brain of an adult individual can be stained with anti-nestin antibodies. Detection of stained cells is an indication of a predisposition to the development of a brain tumor or the presence of a brain  
10 tumor.

#### Brief Description of the Drawings

Figure 1 is the nucleotide and deduced amino acid sequences of the nestin transcript derived from the central nervous system of rat embryos. The three intron  
15 positions (912, 1038 and 1111) are indicated. The canonical poly A addition signal (5924-5929) is underlined.

Figure 2 is the nucleotide sequence of human nestin.

Figure 3 is the deduced amino acid sequence of human  
20 nestin.

#### Detailed Description of the Invention

The present invention relates to a gene of mammalian origin encoding a protein whose expression distinguishes neural multipotential stem cells and brain tumor cells  
25 from the more differentiated neural cell types (e.g., neuronal, glial and muscle cells of the adult brain).

The present invention more particularly relates to the nestin gene and the encoded proteins and their use in



-7-

diagnosing tumors of the brain, such as medulloblastomas, glioblastomas and oligodendroglioma. As described herein, the human nestin gene and the rat nestin gene have been isolated and sequenced. In addition, expression of the nestin gene and its structural features are described. The intracellular distribution of the nestin gene product, both in an embryonic rat brain cell line (ST15A) and human brain tissue, including tumor cells, using immunocytochemical and immunohistochemical techniques is also described. Finally, nestin expression in the developing human central nervous system and brain tumor samples is also described herein.

#### CLONING AND CHARACTERIZATION OF THE RAT AND HUMAN NESTIN GENE

##### 15 The Rat Nestin Gene

A cDNA library in the expression vector lambda gt11 was constructed from poly(A)<sup>+</sup> RNA from CNS of embryonic day 15 (E15) rat embryos as described in detail in Example 1. The library was screened with the monoclonal antibody Rat 401 after induction of protein expression and immunopositive plaques were identified. The cDNA insert from the most immunopositive clone, gt11.401.19, was used as probe to isolate additional clones, gt10.401:16, gt10.401:18 and gt10.401:9, in a lambda gt10 library from rat E15 CNS constructed in parallel with the gt11 library. A number of hybridizing clones were identified and sequenced and together represent 5333 base pair (bp) of transcribed sequence, including 404 bp of 3'

-8-

untranslated sequence and a consensus polyA addition site.

05 The remaining portion of the nestin gene was obtained from genomic sequences. Genomic Southern blots, even when hybridized at low stringency, revealed a pattern consistent with a single rat gene. A genomic rat library in the vector Charon 4A was screened with the nestin cDNA clones. Hybridizing clones were organized by restriction mapping and the region upstream of the cDNA clone gtl0.401:9 was sequenced.

10 An open reading frame continues for an additional 600 bp from the 5' end of the most upstream cDNA clone. This region from the genomic clone Cha.401:14 was subdivided into three different fragments with the restriction enzyme BamH1. These three fragments all hybridized to an mRNA of the correct size, 6.2 kb, when used as probes on Northern blots with rat E15 CNS RNA.

15 The exact location of the 5' end was determined by a combination of S1 nuclease mapping and primer extension. A probe (bp -121 to +194 relative to +1 - the subsequently defined transcription start) was generated overlapping the end of the genomic open reading frame by polymerase chain reaction between two specific primers. This PCR product protected a 194 bp fragment in S1

25 nuclease analysis. Primer extension from the downstream primer used to generate the PCR fragment also produced a 194 bp fragment. The fact that these two experiments gave the same fragment suggests that the transcription start site has been defined.

30 Figure 1 shows the nucleotide sequence of the rat nestin gene. The total length of the nestin gene transcript is 5945 bp, excluding the poly A tail.

-9-

There are two potential initiation codons at positions +127 and +160. Assuming the first methionine is the initiation codon, the coding sequence is 5415 bp. This corresponds to an expected molecular weight of approximately 200 kD; the relative molecular weight as determined by migration on SDS polyacrylamide gels is approximately 240 kD (Hockfield and McKay, supra (1985)). Sequence comparison of cDNA and genomic clones reveals three introns at positions 912, 1038, and 1111 in the transcribed region.

To confirm that the cloned gene indeed represents the gene encoding the Rat401 antigen, a synthetic peptide from the C-terminal 20 amino acids was made. The extreme C-terminus was chosen because it is not included in the clone recognized by the original monoclonal antibody, and therefore represents a distinct epitope. A polyclonal antiserum against the peptide was raised in rabbits and identifies a band in Western blots and a pattern in immunostaining experiments which are indistinguishable from that recognized by the monoclonal antibody Rat 401.

#### Structural Features of the Rat Nestin Gene

The sequence of the entire transcribed region of the rat nestin gene was compared to the Genbank and EMBL databases. The only significant similarities found are to the five classes of intermediate filaments: acidic keratins (class I); basic keratins (class II); desmin, glial fibrillary acidic protein, peripherin, and vimentin (class III); the neurofilament triplet (class IV); and nuclear lamins (class V) (Steinert and Roop, 1988)).

-10-

The similarity between the nestin gene and the genes of the other five classes of intermediate filaments ranges from 16 to 29 % at the amino acid level in a 307 amino acid long region starting close to the N-terminus of the nestin gene, corresponding to the conserved alpha-helical rod or "core" domain of the intermediate filaments. This region of the predicted nestin amino acid sequence also contains a repeated hydrophobic heptad motif characteristic of intermediate filaments. The degree of amino acid similarity is comparable to that found between different classes of intermediate filament, both in degree and in location, i.e., conserved heptad-containing alpha-helical stretches within the core domain are separated by less well conserved, non-heptad spacers (Steinert, P.M. and D.R. Roop, Ann. Rev. Biochem. 57: 593-625 (1988)). The locations of the nestin gene's three introns are not conserved with respect to those of other classes of intermediate filaments.

In the regions outside the conserved rod domain, no strong similarities to other characterized genes were found. The N-terminal domain is only 11 amino acid residues long, shorter than in other intermediate filaments. The large C-terminal domain loosely resembles those of neurofilaments in that it is highly charged, bears glutamate-rich regions, and features a repeat, in this case the 11 amino acid motif S/P-L-E-E/K-E-X-Q-E-S/L-L-R (underlined residues are strongly conserved). There are approximately 35 of these repeats in the region between amino acids 512 and 1050.

The deduced amino acid sequence of nestin suggests that it is a member of the intermediate filament protein

-11-

family. However, its degree of sequence homology to other intermediate filaments in the core domain, its dissimilarity in the head and tail domains, and its different splicing pattern suggest that it defines a new class of intermediate filament protein.

#### The Human Nestin Gene

As described in detail in Example 5, the human nestin gene was isolated using low stringency DNA hybridization of a human genomic bacteriophage lambda library with a rat nestin probe.

Figure 2 shows the nucleotide sequence of the human nestin gene. The total length of the human nestin gene sequence is 4854 bp. Figure 3 shows the deduced amino acid sequence encoded by the human nestin gene which is 1618 amino acid residues in length. The alignment of the deduced amino acid sequences of human nestin with the deduced amino acid sequence of rat nestin showed greater than 75% sequence similarity between the two sequences, with greater than 60% sequence identity.

#### DEVELOPMENTAL EXPRESSION OF THE RAT AND HUMAN NESTIN GENE AND ENCODED PRODUCT

##### Developmental Expression of the Rat Nestin Gene

During development, CNS stem cells differentiate into neurons and glia on a stereotyped schedule; different brain regions become post-mitotic at different times. The nestin protein was originally identified by the Rat 401 antibody, which transiently stains radial

-12-

glial cells of the neural tube in cross sections of rat embryos (Fredericksen, K. and R. McKay, J. Neurosci 5: 3310-3328 (1988); Hockfield and McKay, supra (1985)).

05 The distribution of staining in the neural tube at different developmental stages, combined with quantitative double-label experiments using tritiated thymidine autoradiography and FACS cell cycle analysis, showed that expression of the Rat 401 antigen was localized to proliferating CNS stem cells during embryogenesis  
10 (Frederiksen and McKay, supra (1988)) and U.S. Patent Applications Serial Nos. 180,548; 201,762 and 603,803 herein incorporated by reference. These experiments have been extended by analyzing the developmental expression of the nestin gene.

15 Northern blots of RNA from E15 rat CNS or from the Rat 401 positive cell line ST15A were probed with nestin cDNA inserts, a single 6.2 kb mRNA species was identified. To investigate the temporal profile of nestin RNA expression in the developing CNS, representatives of  
20 early (cerebrum) and late (cerebellum) developing regions of the brain were compared.

In cerebrum, nestin expression was found to decline from a maximum on day E16, such that only extremely low levels of RNA could be detected at day P9 and P12 and  
25 none at all in the adult cerebral cortex, even after long exposures. The loss of nestin mRNA precisely parallels the decline of Rat 401 positive stem cells in cerebrum during neurogenesis (Frederiksen and McKay, supra (1988)). The postnatal cerebellum expresses the nestin  
30 gene with a maximum on day P5; no expression can be

-13-

detected after day P9 . Again, this pattern closely matches the numbers of Rat 401 positive cells in developing cerebellum.

05       - Nestin is not detected by Northern blot analysis in  
adult tissue. The loss of nestin expression coincides  
with terminal differentiation of these early multi-  
potential cells. There are two important implications of  
these results. First, nestin appears to be a general  
10       marker for the CNS stem cell from the newly closed E11  
neural tube through the postnatally developing  
cerebellum. Second, nestin expression is a consequence  
of transcriptional regulation of the gene.

15       The only expression of nestin observed outside  
developing CNS is traces of nestin message from day P4  
developing skeletal muscle. This finding is consistent  
with previous observations from cross sections of rat  
embryos, where Rat 401 staining in the developing somites  
was observed (Hockfield and McKay, supra (1985)).  
Developing nervous tissue and developing skeletal muscle  
20       may share unexpected kinship with regard to their  
potential fates.

#### Intracellular Distribution of the Nestin Gene Product in the ST15A Cell Line

25       The associations of nestin with different components  
of the cytoskeleton were analyzed by a series of double-  
label immunocytochemistry experiments. These experiments  
were carried out in the immortalized CNS precursor cell  
line ST15A which expresses, in addition to components of  
microtubules and microfilaments, nestin and the inter-

-14-

mediate filament vimentin. ST15A produces a nestin mRNA indistinguishable by Northern blot analysis from that found in E15 CNS.

05 In the first set of double-label experiments the intracellular distribution of the nestin gene product, detected by the monoclonal Rat.401 antibody and an appropriate second antibody, was compared to the distribution of actin, tubulin and vimentin. The latter two  
10 proteins were detected by polyclonal antibodies with matching second antibodies while the actin pattern in microfilaments was visualized by rhodamine-conjugated phalloidin.

The intracellular distributions of nestin, microtubules (tubulin), and microfilaments (actin) are quite  
15 different. Microtubules are more evenly distributed in the cell with no particular aggregation around the nucleus, whereas nestin is found in a fiber-like pattern, with a pronounced perinuclear accumulation. We also observe distinct differences when microfilaments and  
20 nestin are stained in the same cell: microfilaments form straight and parallel fibers, in contrast to the gently curved fibers of nestin.

Differences in nestin and vimentin staining in the same ST15A cells are much less clear. Vimentin staining  
25 is stronger than nestin in the cell periphery, and there are regions where intensity varies between the two proteins, but common features are also evident. The overall organization of the two networks is quite similar, with filament arrays radiating out from a  
30 perinuclear center. Moreover, there are regions in cells where it seems likely that nestin and vimentin colocalize.



-15-

In another set of experiments, microfilaments and microtubules were disrupted by cytochalasin B and colchicine, respectively, and the effects on nestin distribution analyzed. When ST15A cells were treated with 10 ug/ml cytochalasin B for one hour the microfilaments became highly disorganized, while the overall morphology of the nestin network was preserved. Microtubules also remained intact after the cytochalasin B treatment. Colchicine treatment (25 ug/ml) for 24 hours caused a collapse of the microtubules. The nestin network could still be identified, although the perinuclear distribution became much more prominent. A similar pattern was observed for vimentin as has been previously reported (Hynes, R.O. and A.T.Destree, Cell 13:151-163 (1978): Monteiro, M.J. and D.W. Cleveland, J. Cell Biol. 108: 579-593 (1989). Microfilaments were essentially unaffected by the colchicine treatment. The inferred relationship of nestin to intermediate filaments based on its deduced amino acid sequence is thus supported by the intracellular distribution of the nestin gene product under various conditions.

ST15A Cells have Features Expected of a Neural Precursor to Brain Tumors

"The prevailing name 'medulloblastoma' given by Bailey and Cushing (1925), is unfortunate because there is no embryonal cell that has been identified as a medulloblast." This sentence begins the discussion of medulloblastoma in a prominent text on neural tumors (Russell, D. and L.J.Rubinstein, Pathology of Tumors of

-16-

the Nervous System, Baltimore, Williams and Wilkins (1989)). The following presents evidence that a cell line derived from rat cerebellum, ST15A, has features expected of a medulloblast.

05 ST15A was derived from postnatal day 4 rat cerebellum. In the rat, as in other mammals, much of cerebellar development occurs after birth, so that there are a large number of dividing precursor cells present at the time from which this line was derived. Thus, ST15A is  
10 derived from the appropriate tissue and stage of development. It has previously been shown that ST15A cells can differentiate along neuronal and glial pathways. The following experiments, which are described in detail in Example 3, show that ST15A cells can also differentiate  
15 into muscle. This result suggests that a single neuroectodermal cell can give rise to the different cell types found in brain tumors.

The cerebellar cell line ST15A was one of several cell lines obtained by infecting primary rat cerebellar  
20 cells with a recombinant retrovirus carrying SV40 T-antigen. A temperature sensitive allele of T-antigen was used to establish these cell lines. At the permissive temperature (i.e., when the T-antigen protein is active) the cells proliferate and express the intermediate  
25 filament protein nestin.

At the non-permissive temperature the T-antigen protein is rapidly degraded and the cells differentiate into either neurons or glial cells depending on the conditions. The differentiation into neurons and glial  
30 cells is best accomplished by growing ST15A cells in

-17-

co-culture with primary cerebellar cells. Partial differentiation has also been observed when ST15A cells are cultured alone; serum-free medium promotes neuronal differentiation and medium containing fetal calf serum promotes glial differentiation.

ST15A cells also can differentiate into muscle cells. This fate was first noted when spontaneously contracting cells were observed in long term cultures of ST15A at 39°C, the non-permissive temperature for T-antigen. Horse serum is often used to support the differentiation of primary myoblasts. When ST15A cells are grown in horse serum at 39°C, they reproducibly differentiate into multi-nucleate cells which express muscle specific proteins, regenerative action potentials and spontaneously contract.

ST15A cells grown for two days at 33°C in 10% fetal calf serum (FCS) in DMEM grew in a disorganized manner; in contrast ST15A cells grown at 33°C in 10% horse serum at 39°C became aligned. The monoclonal antibody Rat 401 recognizes the 220 kD intermediate filament protein nestin. ST15A cells express nestin at both 33°C and 39°C. A second monoclonal antibody which recognizes a skeletal and cardiac muscle isotype of troponin T was reacted with the cells (Lin, J. Biol. Chem. 263: 7309 (1988)). Immunofluorescence shows that troponin T expression is induced when ST15A cells are cultured at 39°C in horse serum.

Immunoblots of proteins extracted from ST15A cells verify that anti-troponin T antibody recognizes a strong band of 38 Kd, the appropriate molecular weight, in the

-18-

cells grown at 39°C in horse serum; at 33°C no band is present. The troponin band in ST15A co-migrates with troponin extracted from rat postnatal day 4 muscle. The double band seen in both muscle cells and differentiated ST15A cells may represent the adult and embryonic isoforms of troponin T which differ by 3,000 daltons (Lin, J. Biol. Chem. 263: 7309 (1988)). These two isoforms are derived from the same primary transcript by differential splicing. Lanes carrying the same amount of protein were also probed for nestin with the monoclonal antibody Rat 401. Nestin is present at both temperatures but at elevated levels at the higher temperature. The lower bands are proteolytic fragments of nestin which are often present in protein extracts even in the presence of protease inhibitors. Nestin is also present in postnatal rat muscle.

A family of DNA binding factors has been shown to play a critical role in the differentiation of muscle. These proteins, myoD1, myogenin, myf5 and herculin/MRF4, contain DNA binding motifs of the helix-loop-helix type in a region of homology to c-myc (Wright, W.E., et al., Cell. 56: 607 (1989); Miner, J.H. and B.Wold, Proc. Nat. Acad. Sci. 87: 1089-1093 (1990); Davis, R.L. et al., Cell. 51: 987-1000 (1987); Braun, T et al., EMBO J 8: 701-709 (1989); Rhodes S.J. and S.F.Konieczny, genes and dev. 3: 2050-2061 (1989). The expression of these factors at the mRNA level in ST15A cells as they differentiate into muscle was examined.

Northern blot analysis of total RNA from differentiating ST15A cells probed with the indicated sequences was performed. Each probe hybridized only to the bands

-19-

of the expected mobilities. MyoD1 is expressed in passaging ST15A cells at the permissive temperature for the T antigen and continues to be expressed when the cells are shifted to 39°C. Myogenin is not expressed under normal growth conditions but is expressed by 2 days after confluent cells are shifted to 39°C in horse serum (HS); if confluent cells are placed in HS and held at 33°C, the induction of myogenin expression is dramatically inhibited.

Herculin/MRF4 was found in adult skeletal muscle but was not detected in ST15A. Myf5 gave a very weak signal in ST15A cells compared to the myoblast cell line C2C12. Nestin mRNA accumulates with time at 39°C, as does the protein.  $\beta$ -actin serves as a control for the amount of RNA at 33°C and in the short time points at 39°C; the decrease in  $\beta$ -actin signal at 3 and 7 days at 39°C may be due to the changing composition of the cytoskeleton accompanying differentiation, as levels of myoD1 mRNA appear constant throughout the experiment. These results indicate the ST15A cells express myogenic transcription factors and that the induction of myogenin correlates with the differentiation into the muscle fate.

As expected of muscle cells, ST15A cells become multinucleate as they differentiate. The most striking similarity to muscle, which initiated this series of experiments, was that ST15A cells grown to confluence in FCS at 33°C and cultured for up to 15 days at 39°C in 10% horse serum can contract spontaneously. The initial action potentials were generated by a series of hyperpolarizing pulses. The final hyperpolarizing pulse was followed by a train of spontaneous action potentials. Spontaneous action potentials were seen after 12 days at 39°C in 10% horse serum.

-20-

Coincident with the appearance of contracting cells, the resting membrane potential falls and it becomes easier to stimulate trains of action potentials with hyperpolarizing current pulses. ST15A cells also  
05 twitched in response to stimuli which elicited action potentials. These results show that ST15A cells differentiate into muscle, expressing myogenic transcription factors, muscle-specific cytoskeletal components and electrical excitability.

10 An interesting question raised by the properties of ST15A cells is whether primary cerebellar cells can differentiate into the muscle fate. Dissociated cells from postnatal day 5 rat cerebellum were cultured for 6 days under conditions which promote muscle different-  
15 iation of ST15A cells (39°C in the presence of 10% horse serum) and stained with monoclonal antibody against troponin T. These culture conditions promote extensive differentiation of neurons and glia. Troponin T positive cells with an elongated, multinucleate morphology of  
20 muscle cells were also seen, but these cells were very rare, occurring less than one per hundred thousand cells.

#### Developmental expression of Human Nestin Gene

To characterize human nestin and determine the timing of its induction in the normal human developing  
25 CNS, studies were carried out, to examine this class VI IF protein in the normal human developing central nervous system (CNS), human brain tumor derived cell lines, and tissue samples of human CNS tumors.

30 Human nestin exhibited biochemical and immunochemical properties similar to those of rat nestin.

-21-

Further, in the human, nestin was detected immuno-histochemically in several different types of immature human CNS cells, i.e. germinal matrix cells, neuro-epithelial cells lining the central canal, radial glia and endothelial cells. Nestin appeared in these cells at the earliest gestational age (i.e., 6 weeks) examined here and then it declined in all but the endothelial cells at later embryonic stages. Nestin also was detected by immunocytochemistry in 6 of 7 primitive neuroectodermal tumor cell lines and in both of the malignant glioma cell lines examined. In these cell lines, nestin co-localized incompletely with bundles of IFs containing other IF proteins (i.e. vimentin, glial filament, neurofilament). Nestin was ubiquitous in a wide variety of brain tumors, but was most prominent in gliomas. These studies document the existence of a human counterpart of rat nestin.

Distribution of the Nestin Gene Product in the Human Developing CNS

Cervical levels of six developing human spinal cords with gestational ages (GAs) of 6 to 40 weeks were immuno-histochemically stained with anti-nestin 129. The spinal cord of the 6 weeks GA human fetus was composed of three distinct layers, i.e. a layer of primitive neuro-epithelial cells lining the central canal, a mantle layer and an outer marginal layer. Anti-nestin 129 stained most cells in the primitive neuroepithelial layer, i.e. the presumptive multipotential stem cells that give rise to CNS neurons and glia. In addition, thin elongated radial glial fibers extending from the primitive neuro-

-22-

epithelial layer to the subpial region were stained by this antiserum. These neuroepithelial stem cells and radial glial fibers also expressed vimentin but not GFAP. Neuroblasts in the mantle layer (identified by their morphology and NF protein positively) did not express nestin.

At 11 weeks GA, cells in the ependymal layer were positively stained with anti-nestin 129, as were radial glial fibers. These radial glial fibers were also stained with the anti-vimentin MAb. Nestin immunoreactivity decreased at 17 weeks GA and disappeared at 20 weeks GA. From this gestational age to maturity, GFAP positive glial cells and NF positive neurons were seen in the spinal cord, but no elements other than endothelial cells were nestin positive.

The telencephalic germinal matrix of two cases with GAs of 17 and 20 weeks were examined with anti-nestin 129. At both time points, many presumptive neuroepithelial stem cells in the germinal matrix cells were nestin positive. These neuroepithelial precursor cells also were labeled by the anti-vimentin MAb, but no germinal matrix cells were stained by the MABs to GFAP or NF proteins. At 40 weeks GA, a well defined germinal matrix was not evident, but clusters of morphologically immature cells (presumably residual, multipotential precursor cells) were seen in the subependymal region. Nestin immunoreactivity was recognized in only a few of these cells.

To determine if neuroepithelial precursors in the cerebellum also expressed nestin, four developing human cerebelli at GAs of 17 to 40 weeks were studied. Late in



-23-

development, the cerebellar cortex has four distinct layers, i.e. the internal and external granular layers, the Purkinje cell layer and the molecular layer. However, these layers were not yet evident at 17 weeks GA and Purkinje cells were not recognizable by morphological criteria. Nevertheless, immature NF positive cells were observed at 17 weeks GA in the superficial internal granular layer, and these cells probably correspond to nascent Purkinje cells. Nestin immunoreactivity was detected at this time in radial glial cells in the internal granular layer and in radial glial processes that extended to the external granular layer. These radial glial cells, which mature into the Bergmann glia of the adult cerebellum, also were stained by the anti-vimentin MAb (data not shown), but not by the anti-GFAP MAb. However, the anti-GFAP MAb did stain ependymal cells lining the fourth ventricle. At 20 weeks GA, nestin immunoreactivity in radial glia was markedly diminished and by 40 weeks GA, it was weakly present in only a few radial glia fibers, while the blood vessel endothelial cells at 40 weeks GA were clearly nestin positive. In contrast, radial glial fibers expressed both GFAP and vimentin (data not shown) at these time points.

#### 25 Nestin Expression in Brain Tumor-derived Cell Lines

Indirect immunofluorescence studies of 9 well characterized human brain tumor derived cell lines showed variable nestin immunoreactivity in 6 or 7 primitive neuroectodermal tumor (PNET) cell lines, and very intense nestin staining in 2 glioma cell lines. Table 1

-24-

summarizes the data on the expression of nestin and other IF proteins in these cell lines.

TABLE 1

05 IMMUNOFLUORESCENCE ASSAY OF BRAIN  
TUMOR-DERIVED CELL LINES

CELL LINES	NESTIN	VIMENTIN	GFAP	NF-L	NF-M	NF-H
Daoy	-	+	-	-	-	-
D283 Med	+	+	-	+	+	+
D341 Med	+	+	-	-	+	+
D384 Med	+	+	-	+	+	+
D425 Med	+	+	-	+	+	+
D458 Med	+	+	-	+	+	+
CHOP707m	+	+	-	+	-	-
U251 MG	+	+	+	-	-	-

15 Summary of the data obtained from the indirect immunofluorescence studies of each cell line. The patterns of staining are described as - = no staining, + = positive staining. NF-L, NF-M and NF-H are the low, middle and high molecular weight neurofilament subunits, respectively. The following  
20 antibodies were used to obtain these data; nestin (rabbit anti-nestin antiserum 129), vimentin (V9), GFAP (2.2B10), NF-L (NR-4, RMS12), NF-M (RMd020, RM0254, H014), NF-H (DP1, TA51, RM024).

25 Except for CHOP707m, which was derived from a cerebral neuroblastoma (Baker, D.L. et al., Am. Neurol. 28:136 (1990)), the other 6 PNET cell lines were obtained from cerebellar medulloblastomas (see Hockfield, S. et

-25-

al., J. Neurosci 5:3310 (1985)) and citations therein). Daoy does not exhibit any evidence of glial or neuronal differentiation and it was nestin negative, while the other 6 PNET cell lines resembled that of embryonic neuroblasts (He, X. et al., J. Neuropathol. Exp. Neurol. 48:48 (1989)). Double immunofluorescent staining of D283 Med, which is the most differentiated PNET cell line, showed incomplete co-localization of nestin and NF proteins in bundles of IFs in the same cells. Similar observations were made using nestin and vimentin antibodies. CHOP707m expressed extensive nestin immunoreactivity and double immunofluorescence of nestin and vimentin in this cell line co-localized both proteins in the same cells, but the nestin positive filament bundles were located mainly in the perinuclear area while vimentin positive filament bundles were present throughout these cells. The Daoy line expresses vimentin, but not NF proteins, GFAP or other molecular markers of neurons or glia (He, X. et al., J. Neuropathol. Exp. Neurol. 48:48 (1989)). Hence, Daoy is the least differentiated PNET cell line, and it did not express nestin. The foregoing co-localization studies also indicated that the anti-nestin 129 did not cross react with vimentin or NF proteins.

Since nestin is expressed primarily in CNS stem cells and is eliminated during the progressive maturation of the progeny of these stem cells, the Daoy cell line may resemble a CNS precursor that subsequently develops into a vimentin and nestin positive CNS stem cell like those observed here in the human spinal cord at a GA of 6 weeks. Alternatively, tumor cell lines contain genetic

-26-

05 mutations and their phenotype may not fully replicate that of normal cells. However, the 2 glioma cell lines (U251 MG and U373 MG) resembled immature spinal cord radial glia (i.e. at 11 weeks GA) since these cells co-expressed nestin, GFAP and vimentin in the triple fluorescence studies.

10 Finally, Western blot studies were performed which showed that anti-nestin 129 labeled a single band in cytoskeletal extracts of U251 MG cells, and this immuno-  
15 band had a  $M_r$  similar to that of the bacterial TrpE-nestin fusion protein and lower than that of rat nestin expressed in P6 rat cerebellum. The significance of the difference in the  $M_r$  of rat and human nestin is unclear and will require further studies. Since these extracts  
15 contained vimentin, GFAP and other cytoskeletal proteins, these data further demonstrate the specificity of the anti-nestin antiserum for human nestin.

#### THE EXPRESSION OF NESTIN IN BRAIN TUMOR TISSUE

##### Brain Tumors Contain Nestin Positive Cells

20 ST15A cells express properties expected of the precursor to brain tumor cells as described in Example 3. Therefore, brain tumor cells were tested to determine whether they express features in common with ST15A, such as expression of nestin. As described in detail in  
25 Example 6, tissue from five medulloblastomas was obtained. Three of these samples came from recurrent tumors occurring in a 16 year old male, a 26 year old male and a 5 year old female. The remaining tumors were

-27-

newly diagnosed in an 11 year old male and a 10 year old female. The primary tumor in the 11 year old male was also used as a source of cultured cells.

05 Immunohistochemistry on sections of tissue from each  
of the five tumors studied showed regions with large  
numbers of nestin immunoreactive cells. The expression  
of nestin in medulloblastoma tissue was not uniform and  
in some cases parts of the tumor tissue showed a few  
nestin-positive cells among nestin-negative cells. The  
10 pre-immune serum showed no reaction with either human  
embryonic or tumor tissue. The nestin positive cells  
were small with a cytoplasmic distribution of the  
antigen. In one tumor, there were groups of aligned  
nestin positive cells which were very similar in their  
15 morphology to the embryonic neuroepithelium.

A cell line which was generated by growing cells out  
of the medulloblastoma from the 11 year old male was also  
tested for expression of nestin. Immunocytochemistry  
with polyclonal anti-nestin antiserum shows that these  
20 cells are uniformly positive for nestin expression and  
reveals the filamentous pattern characteristic of this  
intermediate filament protein. Western blotting using  
anti-nestin serum shows that the antigen detected in  
these cells comigrates with rat nestin.

25 Sections of normal fetal and adult human cerebellum  
were also stained with polyclonal antiserum against  
nestin. The fetal tissue showed many positive cells in  
the proliferative, external granular layer of the  
developing cerebellum and in the outer part of the  
30 internal granular layer. In contrast, there was no

-28-

immunoreactivity in cerebellum from a 2.5 year old. These results are consistent with the stem cell specific expression of nestin seen in developing mouse and rat brain.

05 Utility

10 The adult cell types of the brain express distinct intermediate filament proteins: neurofilaments are expressed by neurons, peripherin by a subset of neurons (Leonard, Gorham, et al., (1988)), and glial fibrillary acidic protein by astrocytes. The expression of nestin defines an earlier stage in the pathway of intermediate filament gene expression. Early embryonic cells are nestin negative but express certain cytokeratins (Jackson, B.W., C. Grund et al., Differentiation 20: 15 203-216 (1980)). During neurulation neurectodermal cells become nestin positive. Upon terminal differentiation, CNS cells stop producing nestin and express class III or IV intermediate filaments typical of their differentiated cell type.

20 Nestin positive neurectodermal cells have been shown to be capable of differentiating into neuronal, glial and muscle cells (i.e., the different cell types found in tumors). Nestin expression has also been directly detected in tumor tissue and in a medulloblastoma derived cell line. Nestin positive cells derived from tumors are 25 likely to be renewing stem cells which give rise to the differentiated cells in the tumor.

30 The detection of nestin expression in brain cells obtained from an adult using known methods is indicative of the presence of a brain tumor or of neural precursor

-29-

cells capable of developing into a brain tumor. For the purposes of the subject invention, an "adult" is an individual whose central nervous system is developed and, therefore, contains differentiated cells. For example, the central nervous system of human neonates is in continual development up until the age of about one year. Therefore, a human adult, for the purposes of the subject invention, is an individual who is older than about one year.

Nestin mRNA expression in a brain tissue sample can be detected using a DNA probe; this is indicative of expression of nestin, which is, in turn an indication of a predisposition to the development of a brain tumor or of the presence of a brain tumor in the individual. For example, a sample of brain tissue from of an individual can be hybridized to a DNA probe which is complementary to all or a portion of the nestin gene. Detection of hybridization is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor.

Alternatively, expression of the nestin protein can be detected using polyclonal antibodies, (e.g., anti-nestin antiserum 129) or monoclonal antibodies (e.g., Rat 401). For example, cerebral spinal fluid or a serum sample from the brain of an individual can be stained with anti-nestin antibodies. Detection of stained cells is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor.

The finding that neural precursors to tumors express nestin also has important therapeutic implications. For example, the nestin marker enables further investigation

-30-

into the proliferation and differentiation of the stem cells of a brain tumor, so that existing chemotherapies can be implemented at an earlier stage in the tumor development and so that new methods of inhibiting tumor development can be developed.

The present invention will now be illustrated by the following examples, which are not to be seen as limiting in any way.

#### EXEMPLIFICATION

##### 10 EXAMPLE 1 Cloning and Characterization of the Rat Nestin Gene

###### Construction and Screening of Libraries

15  $\lambda$ gt10 and  $\lambda$ gt11 cDNA libraries were constructed from oligo-dT-primed poly(A)+ RNA from embryonic day 15 dissected rat CNS according to procedures which were slightly modified from those described (Young, R.A. and R.W.Davis, Proc. Natl. Acad. Sci. USA. 80: 1194-1198 (1983)). The second strand cDNA was synthesized using the RNase H modification of the Okayama-Berg method

20 (Gubler, U. and R. Hoffman, Gene 25: 263-269 (1983)). After induced protein expression, the  $\lambda$ gt11 library was screened with the monoclonal antibody Rat 401 (undiluted hybridoma supernatant) and positive clones were visualized with an anti-mouse second antibody conjugated to

25 alkaline phosphatase (Promega Protoblot). Immunopositive clones were selected and the cDNA insert from the clone giving the strongest hybridization signal,  $\lambda$ gt11.401:16,



-31-

was  $^{32}\text{P}$ -labelled (Feinberg, A.P. and B. Vogelstein, Anal. Biochem. **137**: 266-267 (1983) and used as probe to screen the  $\lambda$  gt10 library (Benton W.D. and R.W. Davis, Science. **196**: 180-182 (1977); Thomas, P.S. Proc. Natl. Acad. Sci. USA. **77**: 5201-5205 (1980)). Several clones were identified. A commercial rat genomic DNA library (Clontech) in the vector Charon 4A was screened with  $^{32}\text{P}$ -labelled nestin cDNA inserts as above. A number of hybridizing clones were obtained, and a genomic map comprising 20 kb of the nestin gene region was established by restriction mapping of the genomic clones, cross-hybridizations, and sequencing of selected genomic subclones.

#### DNA Sequencing and Computer Analysis

cDNA inserts and genomic subclones were sequenced after subcloning into M13mp18, M13mp19, or Bluescript KS+ (Stratagene). M13 phage particles and Bluescript were grown and circular DNA prepared according to standard procedures. Both single stranded (M13) and double stranded (Bluescript) sequencing was performed using a modified T7 DNA polymerase (Sequenase, USB), as suggested by the manufacturer. Approximately 90% of the gene, including all exons and cDNA inserts, was sequenced on both strands. Ambiguous regions were further resolved using deoxyinosine in the standard Sequenase protocol. Computer analysis was performed using the University of Wisconsin Genetics Computer Group program package, including the FASTP and TFASTA algorithms (Devereux, Haeberli et al., 1984)). The complete sequence, including introns, will be stored in the EMBL Database.

- 32 -

### Isolation of RNA and Northern Blotting

Tissue was rapidly dissected and quickly frozen in liquid nitrogen. After homogenization in a guanidinium thiocyanate containing buffer, total RNA was isolated by centrifugation through a CsCl cushion (Chirgwin, Przybyla et al., 1979)). Poly(A)+ RNA was isolated by an oligo-dT chromatography column (Aviv and Leder, 1972) or by a poly(A)+ RNA extraction kit (Fast Trak, Invitrogen). For Northern blot analysis RNA was denatured and electrophoresed in 1% agarose gels containing 2.2 M formaldehyde in 1X MOPS buffer (Maniatis, Fritsch et al., 1982) at +4°C for 8 hours. The RNA was then transferred to nitrocellulose filters (GeneScreen Plus, New England Nuclear) (Thomas, 1980)). The filters were prehybridized and hybridized (50% formamide, 0.75M NaCl, 100 mM Tris pH 7.8, 5 mM EDTA, 1% sodium dodecyl sulphate at 42°C for 16 hours) with  $2 \times 10^6$  cpm/ml of the  $^{32}$ P-labeled nestin cDNA insert from  $\lambda$  gt10.401:16 (specific activity  $3 \times 10^8$  cpm/ug). After hybridization, the filters were washed (final wash =  $2 \times 30$  minutes at 65°C in 30 mM NaCl, 4 mM Tris pH 7.8, 0.2 mM EDTA and 0.25% SDS) and exposed to x-ray film with intensifying screens. Quantity and integrity of the RNA were monitored by rehybridizing the blots with a cDNA probe from the human beta-tubulin gene under identical conditions. The size of the hybridizing RNA was determined by running RNA molecules of known size (HMW RNA ladder, BRL) in parallel.

### S1 Nuclease Mapping and Primer Extension

To generate a probe for S1 nuclease protection experiments, we amplified a region of a genomic clone

-33-

between two synthetic primers using the polymerase chain reaction (kit from Perkin-Elmer Cetus). The resulting fragment was 315 bp long and located between bp -121 and +194 (bp +1 indicates the subsequently defined initiation of transcription). After  $^{32}\text{P}$  end-labelling, 300 fmole of the fragment was hybridized to 50 ug of E15 CNS rat total RNA or 50 ug of yeast total RNA according to standard procedures, and the S1 resistant hybrids separated on denaturing polyacrylamide gels (Maniatis, T., E.F. Fritsch et al., Molecular cloning: A laboratory manual (Cold Spring Harbor, New York: (1982)).  $^{32}\text{P}$  end-labeled restriction digested plasmids and sequencing reactions of known DNAs were used as size standards. The dried gels were exposed to x-ray film. Primer extensions were performed by hybridizing 125 fmole of one of the oligonucleotides used in the polymerase chain reaction (pos +170 - +194) to 10 ug of poly(A)+ RNA from rat E15 CNS according to standard procedures. The reaction product was subjected to gel electrophoresis as in the S1 nuclease experiments.

#### Immunocytochemistry

For all experiments cells from the immortalized cell line ST15A were grown for 2 days (to 40% confluence) in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum on sterile, polyornithine coated glass cover slips, as described in U.S. Patent Applications Serial Nos. 07/603,803; 07/201,762 and 180,548. All antibody incubations lasted 60 minutes at room temperature and were followed by three rinses in PBS (0.14M)

-34-

NaCl, 2.7 mM KCl, 8 mM  $\text{Na}_2\text{HPO}_4$  and 8 mM  $\text{KH}_2\text{PO}_4$  pH 7.2). In the double-label experiments with nestin and tubulin antibodies the ST15A cells were briefly rinsed in PBS, presoaked in microtubule stabilizing buffer (MTSB - 2M glycerol, 0.1 M PIPES pH 6.9, 1 mM  $\text{MgSO}_4$ , 2 mM EDTA) for 2 minutes and then fixed for 5 minutes in 4% para-formaldehyde, 0.5% NP-40 in MTSB for 5 minutes. After two brief rinses in PBS and 5 minutes incubation in blocking solution (0.75% Triton X-100, 500 ug/ml bovine serum albumin, 1% horse serum and 0.05% sodium azide), the cells were incubated with a polyclonal rabbit-anti-tubulin antiserum (kind gift of Dr. Frank Solomon) diluted 1:10000 in PBS. This was followed by an incubation with a fluorescein-conjugated goat-anti-rabbit IgG antiserum (Cappel, No. 1612-0081) diluted 1:100 in PBS. The cells were then postfixed in 95% ethanol/5% acetic acid for 5 seconds and rinsed three times in PBS. This was followed by an incubation with undiluted hybridoma supernatant from the monoclonal mouse-anti-nestin antibody (Rat 401). Finally, the cells were incubated with a rhodamine-conjugated polyclonal goat-anti-mouse IgG antibody (1:100 dilution, Cappel No. 2611-0231). After two brief rinses in water, the cells were covered with Immumount (Shandon) and inverted onto clean glass microscope slides. As control experiments for potential cross-hybridization and spillover between the fluorescein and rhodamine spectra in the fluorescence microscope, the first antibodies (rabbit-anti-tubulin and mouse-anti-nestin) were omitted in two separate experiments under otherwise identical conditions. No cross-hybridization

-35-

or significant spillover was observed. In the colchicine experiments the ST15A cells were grown as above and treated with colchicine at 25 ug/ml for 24 hours followed by fixation and staining as described here.

05 In the actin/nestin double-label experiments ST15A cells were grown as above and fixed in 4% paraformaldehyde, 0.1% Triton X-100 in PBS for 5 minutes. After blocking as described above, the cells were incubated with 0.16 uM rhodamine-conjugated phalloidin, as  
10 suggested by the manufacturer (Molecular Probes). Rinses, postfixation and staining with the nestin monoclonal antibody were performed as described above, with the exception that a fluorescein-conjugated sheep-anti-mouse IgG second antibody (diluted 1:100, Cappel No.  
15 1606-3152) was used. No cross-hybridization or spillover was observed after omitting the rhodamin-conjugated phalloidin or the nestin antibody. In the cytochalasin B experiments the ST15A cells were treated with cytochalasin B at a final concentration of 10 ug/ml for 60  
20 minutes, otherwise all immunostaining procedures were identical.

In the vimentin/nestin double-label experiments ST15A cells were fixed in 4% paraformaldehyde, 0.1% Triton X-100 in PBS for 5 minutes. After blocking, the  
25 cells were incubated with a polyclonal goat-anti-vimentin antiserum (dilution 1:20, MILES 65-794) followed by a fluorescein-conjugated rabbit-anti-goat IgG second antibody (dilution 1:100, ICN 65-176). Incubation times and rinsing procedures were as described above. Post-  
30 fixation and incubation with the anti-nestin antibody

-36-

were performed as in the tubulin/nestin experiments. No cross-hybridization or spillover was detected after omitting the anti-vimentin and anti-nestin antibodies, respectively. In all experiments the mounted cells were  
05 photographed on Kodak Ectachrome 400 ASA using a 63X objective on a Zeiss fluorescence microscope.

EXAMPLE 2 Production of the Rat 401 Monoclonal Antibody

Developing monoclonal antibodies were generated to fixed spinal cord from embryonic day 15 (E15) rats.  
10 Timed pregnant female rats were obtained from Taconic Animal Supply Co. and housed until the appropriate gestation age. Uteri were dissected from pregnant animals into ice-cold phosphate buffer (pH 7.4), individual embryos were removed and the spinal cord was  
15 dissected free of other tissue into fixative (4% paraformaldehyde in 0.1 M phosphate buffer).

The immunization and fusion protocol has been described previously (McKay and Hockfield, Proc. Natl. Acad. Sci. USA 79: 6747-6751 (1982)). Briefly, BALB/c  
20 mice received two intraperitoneal immunizations with fixed tissue homogenized in saline and suspended in an equal volume of Freud's complete adjuvant and a final intravenous boost of unfixed tissue with adjuvant. Spleen cells from immunized mice were fused with NS1  
25 myeloma cells and resulting hybrid cell lines were screened immunohistochemically on 100  $\mu$ m-thick Vibratome sections of 4% paraformaldehyde-fixed E15 rats. Lines producing antibodies of interest were cloned in soft agar or by limiting dilution and antibody subclass was deter-  
30 mined by the Ouchterlony method (Ouchterlony and Nilsson,

-37-

Immunodiffusion and Immunelectrophoresis. In Handbook of Experimental Immunology D.M. Weir, ed., Blackwell Scientific Publications, Oxford (1978).

#### Immunohistochemistry

05 Tissue for immunohistochemistry was fixed either by  
intravascular perfusion or by immersion. All postnatal  
material was fixed by intravascular perfusion (under  
ether anesthesia) with 4% paraformaldehyde at pH 7.4 for  
light microscopy or with either 4% paraformaldehyde at pH  
10 10.0 or 4% paraformaldehyde, 0.1% glutaraldehyde at pH  
7.4 for electron microscopy. The CNS was dissected out  
and stored in 0.1 M phosphate buffer (pH 7.4) with 0.1%  
sodium azide. For prenatal material, pregnant females  
were ether anesthetized, uteri were dissected into cold  
15 0.1 M phosphate buffer and the females were killed by  
cervical dislocation. Embryos were dissected indi-  
vidually into Sylgard-coated Petri dishes and pinned in  
place. Intravascular perfusion was performed by opening  
the skin of the thoracic cavity, nicking the right atrium  
20 with a fine forceps and placing a 25 gauge needle into  
the left ventricle. One to 5 ml of fixative (as above)  
were injected into the heart. The embryos were placed in  
a vial containing the same fixative for 4 hr and then  
stored in phosphate buffer. E12 to term embryos were  
25 fixed by perfusion (using a 30 gauge needle for E12  
embryos); E10 and E11 embryos were fixed by immersion in  
fixative. E13 and older animals were sectioned at 50 to  
100  $\mu$ m on a Vibratome without additional support. E10 to  
E12 embryos were embedded in 15% gelatin before  
30 sectioning.

-38-

Immunohistochemical reactions were performed on free-floating sections by sequential incubations in monoclonal antibody as full-strength supernatant (12 to 20 hr); horseradish peroxidase (HRP)-conjugated goat anti-mouse antibody at a dilution of 1:100 in tissue culture medium with 10% serum; and 3,3'-diaminobenzidine (DAB; 0.025%) with  $H_2O_2$  (0.002%). Peroxidase-conjugated lectins (Sigma Chemical Co.) were used at serial dilutions in the same manner with the omission of the peroxidase-conjugated goat anti-mouse antibody. For light microscopy, 2% Triton X-100 was included in both antibody solutions. For electron microscopy sections carried through this procedure (without Triton X-100) were post-fixed in 2%  $O_2O_4$  embedded in Epon/Araldite between plastic coverslips, and cut at 1  $\mu m$  and 0.12  $\mu m$  for correlative light and electron microscopy. For high resolution light microscopy, sections (Triton-treated) were embedded in plastic as for electron microscopy and sectioned at 2  $\mu m$ .

## 20 Immunoblots

The apparent molecular weights of antigens recognized by antibodies were determined using the method of Towbin *et al.*, Proc. Natl. Acad. Sci. USA 76: 4350-4354 (1979)). Proteins were extracted from unfixed tissue with sodium dodecyl sulfate (SDS) and  $\beta$ -mercaptoethanol and run on an SDS-polyacrylamide gel. The proteins were electroblotted onto nitrocellulose paper which was then exposed to primary (1 hr) and secondary (1 hr) antibodies. HRP was visualized with 4-chloronaphthol.



-39-

EXAMPLE 3 Characterization of Rat Nestin from the  
ST15A Cell Line

ST15A cells were derived from rat postnatal day 2 cerebellum after infection of a primary culture with a retrovirus transducing the temperature sensitive, tsA58 allele of SV40 T antigen (Frederiksen et al., 1988)). ST15A cells were plated on tissue culture plastic coated with polyornithine and passaged at the permissive temperature, 33°C, in Dulbecco's modified Eagle's medium (DME) supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate and 15 mM HEPES, pH 7.2, containing 10% fetal calf serum (FCS). For the experiment in Figure 8, confluent cells (labelled 'day 0') were transferred into DME/10% horse serum (HS) and cultured at 33°C or 39°C as marked for the indicated time. 39°C is the non-permissive temperature for tsA58.

Primary cultures were prepared from P4 rat cerebellum as previously described (McKay, R.D.G. et al., Cold Spring Harbor Symp. Quant. Biol. (1990). Briefly, tissue was dissociated by trypsinization followed by trituration. Cells were plated on polyornithine-coated coverslips in DME/10% FCS/10% HS and grown at 39°C for 24 hours. The medium was changed to DME/10% HS and culture continued for 6 days.

The medulloblastoma cell line was derived by growing cells from dispersed, fresh medulloblastoma tissue. The tissue was removed from a previously untreated tumor in a 10 year old female. The cell line has not been cloned. Cells were passaged in Ham's F12 supplemented with 20mm L-glutamine and 10% FCS.

-40-

Immunocytochemistry

Cells for immunocytochemistry were fixed in 4% paraformaldehyde buffered with 50 mM sodium phosphate, pH 7.2, for troponin staining or buffered with 50mM sodium borate, pH 9.5, for nestin staining. The antibodies used were mouse monoclonal Rat 401 (Hockfield and McKay, supra (1985)), mouse monoclonal antibody against troponin T (Sigma T-6277), and rabbit polyclonal serum raised against the C-terminal region of nestin expressed in bacteria. Fixed cells were incubated in primary antibody for one hour, rinsed 3 times in PBS, incubated for one hour with flourescein-conjugated goat anti-rabbit serum (Organon-Tetnika) and mounted in Immumount (Shandon, Inc.) with 2% DABCO. Medulloblastoma tissue and normal human cerebellum were frozen and subsequently fixed in formalin. 50 um sections were cut and stained with rabbit polyclonal serum raised against the C-terminal region of nestin.

Isolation of RNA and Northern blotting

Cultured cells were scraped into 5M guanidine thiocyanate/25 mM citrate, pH 6.8/ 0.5% N-lauryl sarcosine/0.1  $\beta$ -mercaptoethanol and RNA pelleted through a CsCl step gradient (Chirgwin et al., 1979). 20 ug of total RNA was electrophoresed through a 1% agarose/2.2M formaldehyde gel and transferred to nitrocellulose. Prehybridization and hybridization were in 50% formamide/5xSSC/5X Denhardt's/0.1% SDS/0.1 mg/ml salmon sperm DNA at 42°C. Filters were washed in 2xSSC/0.1% SDS at room temperature and in 0.1xSSC/0.1% SDS at 55°C and

-41-

exposed to X-ray film with an intensifying screen. Filters were stripped by washing 6 times for 3 minutes each in boiling 0.05xSSC/0.1% SDS/20 mM EDTA, pH8, and rehybridized. Probes were prepared for hybridization by random priming on fragments isolated on low melting temperature agarose gels (Feinberg and Vogelstein, supra (1984)). The following plasmids were used to generate probes: mouse myoD1 cDNA pEMC11s (Davis, R.L. et al., Cell. 51: 987-1000 (1987); mouse myogenin cDNA p65 (Wright et al., supra (1989); rat nestin cDNA p401-16; human B-actin cDNA (Gunning, P. et al., Mol. Cell. Biol. 3: 787-795 (1983)).

#### Western Blotting

Proteins were extracted from cultured cells or rat postnatal day 5 limb muscle (Sprague-Dawley; Taconic, Inc.) in the presence of aprotinin and PMSF in 2% SDS, 5 mM EDTA and 15% glycerol in 0.1 M Tris, pH 6.8. 50 ug of total protein was run in each lane of an SDS reducing gel according to Laemmli (Laemmli 1970); proteins were transferred to Immobilon P membranes (Millipore) using a Hoeffer transfer apparatus at 50 volts for 4 hours. Filters were blocked in 5% BLOTTO (Johnson, D.A. et al., Gene Anal. Tech. 1: 3-8 (1984)) overnight. Rat 401 hybridoma supernatant was used undiluted; troponin T antibody was diluted 1:500; polyclonal anti-nestin antiserum was diluted 1:1000; HRP conjugated secondary antibodies were purchased from BioRad.

-42-

EXAMPLE 4 Characterization of Human Nestin

To characterize human nestin, a new antiserum, designated anti-nestin 129, was raised to recombinant rat nestin expressed in E. coli since the rat-401 mAb failed to recognize human nestin in immunohistochemical and immunochemical assays.

Production of Anti-nestin Antiserum

The insert from the clone  $\lambda$ gt10 401:16 (Lendahl, U. et al., Cell 60: 585 (1990)) was isolated and ligated into the pATH1 vector (Tzagaloff, A. et al., J. Biol. Chem. 262: 17163 (1986)), producing a clone in which the bacterial TrpE protein (37.2 kD) was fused at its C terminus with the last 1197 amino acids of nestin. The fusion protein was induced in E. coli strain HB101 with 20  $\mu$ g/ml indoleacrylic acid (Sigma I1625) for 4 hours as described in (Tapscott, S. et al., Science 242: 405 (1988)). The TrpE-nestin fusion protein migrates at greater than 200 kD in 6% SDS PAGE gels, making it the largest protein in the bacterial lysate. The fusion protein was purified by cutting the top band from the gel. Anti-nestin antiserum 129 was produced by injecting the fusion protein subcutaneously into a female NZW rabbit. The characterization of anti-nestin 129 and its specificity for human nestin is described here.

First, this antiserum was compared to the Rat-401 MAb in immunochemical and immunohistochemical studies of postnatal day 6 (P6) rat cerebellum. Then, anti-nestin 129 was used to probe human CNS tissues, tumors and tumor derived cell lines. To determine if nestin co-localized with other IF proteins, double and triple immuno-

-43-

fluorescence experiments also were performed using anti-nestin 129 and antibodies to vimentin, glial fibrillary acidic protein (GFAP) and neurofilament (NF) triplet proteins. Finally, Western blots were performed on cytoskeletal extracts of some of the tumor derived cell lines to identify human nestin and to compare it with rat nestin as well as with the class III and IV IF proteins expressed by CNS cells.

#### Specificity of Anti-Nestin 129 in the Rat

The rabbit anti-nestin antiserum produced as described herein, i.e. anti-nestin 129, yielded results identical to those produced by the Rat-401 MAb when both were used to probe P6 and adult rat CNS tissues by immunoblots and immunohistochemistry. On immunoblots of cytoskeletal extracts from P6 rat cerebellum separated by SDS-PAGE, anti-nestin 129 identified a band with an apparent  $M_r$  of 240 kD. This band was identical to the band labeled by Rat-401. Further, both antibodies labeled similar bands in lysates of the induced bacteria producing the TrpE-nestin fusion protein.

In immunohistochemical studies of the P6 rat cerebellum, anti-nestin 129 labelled radial glial cells very intensively. The processes of these cells radiated from the internal granular layer to the external granular layer. These radial glia also were labeled with Rat-401 and the anti-vimentin MAb. Neither of the anti-nestin antibodies stained immature Purkinje cells or other neurons. In contrast to the P6 rat cerebellum, the anti-nestin 129 and Rat-401 antibodies only stained blood vessel endothelial cells in the adult rat cerebellum,

-44-

while the anti-vimentin MAb (V9) labelled blood vessels, white matter astrocytes and radial glial fibers.

Furthermore, nestin immunoreactivity was demonstrated by both anti-nestin antibodies in immature skeletal muscle of the P6 rat as well as in Schwann cells of adult rats (data not shown).

These studies demonstrate that this new anti-nestin antiserum yields immunohistochemical results in the human developing spinal cord, cerebrum and cerebellum that were nearly identical to those produced by Rat-401 and the anti-nestin antiserum in the rat.

#### EXAMPLE 5 Cloning of the Human Nestin Gene

The human nestin gene was isolated using low-stringency DNA hybridization of a human genomic bacteriophage lambda library with a rat nestin probe.

A genomic library in the vector EMBL 3 made from partially Sau3A-digested human genomic DNA (Clontech) was screened using a <sup>32</sup>P-labelled probe derived from the rat nestin gene starting at 202 nucleotides upstream of the translation start and ending at nucleotide 483 after the translation start (Lendahl, et al., 1990) under the following conditions: prehybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 micrograms/ml salmon sperm DNA at 65°C for 1 hour; hybridization in the same conditions except for using 50 micrograms/ml salmon sperm DNA and hybridizing for 16 hours. The filters were then washed for 2 X 30 minutes at 48°C in 2X SSC and 0.25% SDS.

One positive clone out of  $1.5 \times 10^6$  plaques screened contained the entire human nestin gene and was

-45-

chosen for further analysis. This clone was purified and grown in large scale (Maniatis, et al., 1989). A restriction map was established and EcoRI fragments were subcloned in the plasmid vector Bluescript KS I (Stratagene).

05

EXAMPLE 6 Nestin Expression in Brain Tumor Samples

Immunohistochemical data on the expression of nestin, other IF proteins and synaptophysin (SYP) by a diverse group of 34 CNS tumors are summarized in Table 2.

**TABLE 2** Immunoperoxidase staining of brain tumors

No.	Age (yr)	Sex	Material	Location	Nestin	Vim	GFAP	NP	SYP
<b>PNETs</b>									
<b>PNET+N,G</b>									
1	1	M	F	posterior fossa	+	+++	+	+	-
2	1	M	F	posterior fossa	+	+++	+	+++	+++
3	5	F	F	posterior fossa	++	+	+	+	++
4	1	M	F	posterior fossa	++	++	+	+	+++
5	1 month	F	F	posterior fossa	++	++	+	+++	+
6	5	M	F	posterior fossa	++	+++	++	+	+
7	8 months	F	F	posterior fossa	+++	+++	++	+	+
8	1	F	F+P	posterior fossa <sup>d</sup>	++	++	+	+	+++
<b>PNET+N</b>									
9	13	F	F	posterior fossa <sup>b</sup>	-	++++	-	++	++
10	6	M	F	posterior fossa	-	+	E	++	++
<b>PNET+G</b>									
11	11	M	F	posterior fossa	+	+++	++	P	+
12	4	F	F	posterior fossa	+++	+++	++	P	+++
<b>PNET,NOS</b>									
13	2	M	F	pineal	++	++	-	-	++
14	11	M	F	posterior fossa <sup>b</sup>	-	+	-	-	++
15	14	M	F	posterior fossa <sup>b</sup>	+	+++	-	-	+
<b>Astrocytic tumors</b>									
16	2	F	F	posterior fossa <sup>c</sup>	++	++++	++++	-	-
17	6	F	F+P	posterior fossa <sup>c</sup>	-	++++	+++	-	-
18	6 months	M	F	hypothalamus <sup>c</sup>	++	+++	+++	-	-
19	17	M	F+P	posterior fossa <sup>d</sup>	-	+++	+++	-	-
20	3	M	F	frontal <sup>e</sup>	+++	NA	+	-	NA
21	6	M	F+P	occipital <sup>f</sup>	+++	+++	+++	-	-
22	13	M	F	posterior fossa <sup>f</sup>	+++	+++	+++	-	-
<b>Ependymomas</b>									
23	3	F	F	parietal	++	++++	+	-	-
24	2	M	F	posterior fossa	+	-	+	-	-
25	3	M	F+P	posterior fossa	++	++++	+++	-	-
26	5	M	F+P	posterior fossa	++	++++	++	-	-
27	15	F	F+P	cerebral hemisphere <sup>g</sup>	++	++++	+	-	-
28	8	F	F+P	parietal <sup>h</sup>	+++	+++	++	-	-
<b>Choroid plexus papillomas</b>									
29	1	M	F	posterior fossa	-	+++	+	-	-
30	2	M	F	lateral ventricle <sup>i</sup>	-	+++	-	-	-
<b>Gangliogliomas</b>									
31	2	F	F	temporal	++	+++	+++	+	-
32	1	F	F	occipital	++	++	+++	+	-
<b>Meningiomas</b>									
33	8	M	F	posterior fossa	++	++	-	-	-
34	27	M	F+P	anterior fossa	+	+++	-	-	-



Legend to Table 2

Summary of the data obtained from the immunoperoxidase studies of each of the brain tumor samples. Results from frozen (F) and paraffin (P) material are combined in the cases in which both kinds of material were studied. The patterns of staining are described as - = no staining, + = positive in less than 5% of tumor cells, ++ = positive in between 5 - 50% of tumor cells, +++ = positive in between 50 - 95% of tumor cells, ++++ = positive in more than 95% of tumor cells, E = positive cells equivocal, i.e. reactive astrocytes of tumor cells, P = interstitial filamentous staining in the neuropil. The following antibodies were used to obtain these data; nestin (rabbit anti-nestin antiserum 129), vimentin (V9), GFAP (2.2B10), NF (RMS12, RMO20, RMO254, HO14, DP1, TA51, RMO24). NA = not available due to limited material; PNET+N,G = PNET with neuronal and glial differentiation; PNET+N = PNET with neuronal differentiation; PNET + G = PNET with glial differentiation; PNET-NOS = PNET, not otherwise specified. <sup>a</sup>recurrent tumor of case 7, <sup>b</sup>autopsy case, <sup>c</sup>benign astrocytoma <sup>d</sup>genign astrocytoma + angioma, <sup>e</sup>anaplastic astrocytoma <sup>f</sup>glioblastoma multiforme, <sup>g</sup>mixed ependymo-astrocytoma, <sup>h</sup>anaplastic ependymoma, <sup>i</sup>choroid plexus carcinoma.

- 25 The results from frozen and ethanol-fixed paraffin-embedded samples from the same biopsy are combined in the 10 cases from which these paired samples were available. Notably, the immunohistochemical results obtained from each of these paired samples were identical, although the
- 30 primary antibodies were used at slightly higher dilutions in the frozen material compared with the paraffin material.

- 35 PNETs The anti-nestin antiserum stained tumor cells in 12 of the 15 PNETS examined here. Nestin immunoreactivity was found in the cytoplasm of individual neoplastic cells. Reactivity was also observed in the coarse processes of large stellate cells, as well as in

-48-

blood vessel endothelial cells. In the PNETs with an insular architecture, nestin reactive cells were found both within and outside the islands. Since PNETs express SYP and other neuroendocrine markers as well as all classes of IF proteins (Gould, V.E. et al., Human Pathol. 21:245 (1990)), the PNETs were probed for SYP. These studies demonstrated SYP in 14 of 15 PNETs. Vimentin was detected in all the PNETs. Ten of the 15 PNETs expressed GFAP and 10 expressed 1 or more NF proteins. Double immunofluorescence studies of nestin and NF proteins, as well as nestin and GFAP on frozen sections showed their co-localization in the same cells (data not shown). The immunohistochemical data on GFAP and NF proteins suggested that 15 PNETs could be classified into 4 subtypes, i.e. 3 cases of PNET-NOS (not otherwise specified), 2 cases of PNET+G (PNET with glial differentiation), 2 cases of PNET+N (PNET with neuronal differentiation) and 8 cases of PNET+G,N (PNET with glial and neuronal differentiation). All 8 cases of PNET+G,N showed nestin immunoreactivity and 2 cases of PNET+G expressed nestin. On the other hand, 2 cases of PNET+N were negative for nestin. These results may indicate that PNETs with neuronal differentiation tend to lack nestin expression in contrast to PNETs with glial differentiation. This phenomenon paralleled normal differentiation in that postmitotic neuroblasts did not express nestin whereas radial glial cells did, even at relatively late developmental stages. Two of 3 cases of PNET-NOS showed nestin positive tumor cells. This immunohistochemical phenotype might represent primitive neuro-epithelial cells or immature radial glial cells in the

-49-

developing CNS. The other PNET-NOS case was negative for nestin and it might correspond to extremely immature CNS precursor cells just like the most undifferentiated PNET cell line (i.e. Daoy).

05        Astrocytic tumors Five of 7 astrocytic tumors displayed positivity for nestin in the perinuclear cytoplasm of the tumor cells as well as in the cytoplasmic processes of these cells. Additionally blood vessel walls were nestin positive. Two of 4 benign  
10 astrocytomas were negative for nestin. One anaplastic astrocytoma and 2 glioblastomas showed strong nestin positivity. In the glioblastomas, multinuclear giant cells were stained strongly.

Ependymomas Six tumors were histologically  
15 diagnosed as ependymomas and all showed nestin positivity. Immunoreactivity for nestin was found in the tumor cells which formed perivascular pseudo-rosetts. One ependymoma (with marked nuclear pleomorphism, necrosis and a few mitoses) was diagnosed as an ana-  
20 plastic ependymoma and it demonstrated the strongest nestin immunoreactivity among the 6 ependymomas. Expression of nestin described here indicates that nestin is the dominant IF protein of early CNS precursor cells. However, nestin is rapidly extinguished in neurons, glia  
25 and other CNS cell types derived from these precursors. The mechanisms responsible for these rapid changes in IF protein expression are unknown.

          In contrast to the restricted expression of nestin in normal developing CNS cells, this IF protein was  
30 ubiquitously present in a wide variety of CNS neoplasms.

-50-

Nestin was expressed in a subset of PNETs, but was more abundant in gliomas. The presence of nestin in diverse types of neuroepithelial tumors is similar to vimentin; but distinct from the distribution of GFAP and NF

05 proteins in CNS tumors.

-51-

CLAIMS

1. Isolated DNA encoding a protein of mammalian origin whose expression distinguishes neural multipotential stem cells from neuronal, glial and muscle cells.
- 05 2. Isolated DNA of Claim 1 wherein the protein is nestin.
3. Isolated DNA of Claim 2 having the nucleotide sequence of all or a portion of the nucleotide sequence of Figure 1 or Figure 2.
- 10 4. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 1.
5. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 2.
- 15 6. Isolated DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 1.
7. Isolated DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 2.
- 20 8. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 1 which encodes a protein of mammalian origin whose expression distinguishes neural multipotential stem cells from neuronal, glial and muscle cells.
- 25

-52-

9. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 2 which encodes a protein of mammalian origin whose expression distinguishes  
05 - neural multipotential stem cells from neuronal, glial and muscle cells.
10. A DNA probe for the detection of a brain tumor, the probe comprising DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 1  
10 or to all or a portion of a DNA sequence sufficiently homologous to the DNA sequence of Figure 1 that it encodes a protein which is indicative of a predisposition to the development of a brain tumor in an individual.
- 15 11. A DNA probe for the detection of a brain tumor, the probe comprising DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 2 or to all or a portion of a DNA sequence sufficiently homologous to the DNA sequence of Figure 2 that  
20 it encodes a protein which is indicative of a predisposition to the development of a brain tumor in an individual.

-53-

12. A method of diagnosing a predisposition to the development of a brain tumor or the presence of a brain tumor in an individual, comprising:

- 05 a) obtaining a sample from the brain of an individual;
- b) treating the sample in a manner that renders DNA present in the sample available for hybridization with a complementary DNA sequence thereby producing a treated sample;
- 10 c) contacting the treated sample with at least one probe which is a DNA sequence which hybridizes with all or a portion of a DNA sequence represented in Figure 1 or Figure 2; and
- 15 d) detecting the hybridization of DNA from the sample with the DNA probe; wherein hybridization is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor in the individual.

13. A method of diagnosing a predisposition to the development of a brain tumor or the presence of a brain tumor in an individual, comprising:

- 20 a) obtaining a sample from the brain of an individual;
- 25 b) treating the sample in a manner that renders the cells available for staining with a nestin-specific antibody thereby producing a treated sample;
- c) staining the cells of the treated sample with labelled anti-nestin antibodies; and
- 30 d) detecting stained cells as an indication of a predisposition to the development of brain tumors or the presence of a brain tumor in an individual.

-54-

14. A method of Claim 13 wherein the antibodies are labelled with a label selected from the group consisting of fluorescein, rhodamine and peroxidase.
- 05 15. A method of Claim 13 wherein the nestin-specific antibody is a polyclonal antibody.
16. A method of Claim 15 wherein the polyclonal antibody is anti-nestin 129.
- 10 17. A kit for diagnosing a brain tumor, comprising at least one nucleic acid probe, which hybridizes with a DNA sequence represented in Figure 1 or Figure 2.
18. A kit for diagnosing a brain tumor, comprising a first antibody which is nestin-specific and a second antibody which binds to the first antibody and is labelled.
- 15 19. A nestin-specific antibody or a pharmaceutical composition comprising a nestin-specific antibody formulated in a solution for parenteral administration for use in in vivo diagnosis, e.g., tumor imaging.
- 20 20. Use of a nestin-specific antibody for the manufacture of a diagnostic agent for in vivo tumor imaging.



Figure 1

1 120  
22 240  
30 260  
38 280  
46 300  
54 320  
62 340  
70 360  
78 380  
86 400  
94 420  
102 440  
110 460  
118 480  
126 500  
134 520  
142 540  
150 560  
158 580  
166 600  
174 620  
182 640  
190 660  
198 680  
206 700  
214 720  
222 740  
230 760  
238 780  
246 800  
254 820  
262 840  
270 860  
278 880  
286 900  
294 920  
302 940  
310 960  
318 980  
326 1000  
334 1020  
342 1040  
350 1060  
358 1080  
366 1100  
374 1120  
382 1140  
390 1160  
398 1180  
406 1200  
414 1220  
422 1240  
430 1260  
438 1280  
446 1300  
454 1320  
462 1340  
470 1360  
478 1380  
486 1400  
494 1420  
502 1440  
510 1460  
518 1480  
526 1500  
534 1520  
542 1540  
550 1560  
558 1580  
566 1600  
574 1620  
582 1640  
590 1660  
598 1680  
606 1700  
614 1720  
622 1740  
630 1760  
638 1780  
646 1800  
654 1820  
662 1840  
670 1860  
678 1880  
686 1900  
694 1920  
702 1940  
710 1960  
718 1980  
726 2000  
734 2020  
742 2040  
750 2060  
758 2080  
766 2100  
774 2120  
782 2140  
790 2160  
798 2180  
806 2200  
814 2220  
822 2240  
830 2260  
838 2280  
846 2300  
854 2320  
862 2340  
870 2360  
878 2380  
886 2400  
894 2420  
902 2440  
910 2460  
918 2480  
926 2500  
934 2520  
942 2540  
950 2560  
958 2580  
966 2600  
974 2620  
982 2640  
990 2660  
998 2680  
1006 2700  
1014 2720  
1022 2740  
1030 2760  
1038 2780  
1046 2800  
1054 2820  
1062 2840  
1070 2860  
1078 2880  
1086 2900  
1094 2920  
1102 2940  
1110 2960  
1118 2980  
1126 3000  
1134 3020  
1142 3040  
1150 3060  
1158 3080  
1166 3100  
1174 3120  
1182 3140  
1190 3160  
1198 3180  
1206 3200  
1214 3220  
1222 3240  
1230 3260  
1238 3280  
1246 3300  
1254 3320  
1262 3340  
1270 3360  
1278 3380  
1286 3400  
1294 3420  
1302 3440  
1310 3460  
1318 3480  
1326 3500  
1334 3520  
1342 3540  
1350 3560  
1358 3580  
1366 3600  
1374 3620  
1382 3640  
1390 3660  
1398 3680  
1406 3700  
1414 3720  
1422 3740  
1430 3760  
1438 3780  
1446 3800  
1454 3820  
1462 3840  
1470 3860  
1478 3880  
1486 3900  
1494 3920  
1502 3940  
1510 3960  
1518 3980  
1526 4000  
1534 4020  
1542 4040  
1550 4060  
1558 4080  
1566 4100  
1574 4120  
1582 4140  
1590 4160  
1598 4180  
1606 4200  
1614 4220  
1622 4240  
1630 4260  
1638 4280  
1646 4300  
1654 4320  
1662 4340  
1670 4360  
1678 4380  
1686 4400  
1694 4420  
1702 4440  
1710 4460  
1718 4480  
1726 4500  
1734 4520  
1742 4540  
1750 4560  
1758 4580  
1766 4600  
1774 4620  
1782 4640  
1790 4660  
1798 4680  
1806 4700  
1814 4720  
1822 4740  
1830 4760  
1838 4780  
1846 4800  
1854 4820  
1862 4840  
1870 4860  
1878 4880  
1886 4900  
1894 4920  
1902 4940  
1910 4960  
1918 4980  
1926 5000  
1934 5020  
1942 5040  
1950 5060  
1958 5080  
1966 5100  
1974 5120  
1982 5140  
1990 5160  
1998 5180  
2006 5200  
2014 5220  
2022 5240  
2030 5260  
2038 5280  
2046 5300  
2054 5320  
2062 5340  
2070 5360  
2078 5380  
2086 5400  
2094 5420  
2102 5440  
2110 5460  
2118 5480  
2126 5500  
2134 5520  
2142 5540  
2150 5560  
2158 5580  
2166 5600  
2174 5620  
2182 5640  
2190 5660  
2198 5680  
2206 5700  
2214 5720  
2222 5740  
2230 5760  
2238 5780  
2246 5800  
2254 5820  
2262 5840  
2270 5860  
2278 5880  
2286 5900  
2294 5920  
2302 5940  
2310 5960  
2318 5980  
2326 6000  
2334 6020  
2342 6040  
2350 6060  
2358 6080  
2366 6100  
2374 6120  
2382 6140  
2390 6160  
2398 6180  
2406 6200  
2414 6220  
2422 6240  
2430 6260  
2438 6280  
2446 6300  
2454 6320  
2462 6340  
2470 6360  
2478 6380  
2486 6400  
2494 6420  
2502 6440  
2510 6460  
2518 6480  
2526 6500  
2534 6520  
2542 6540  
2550 6560  
2558 6580  
2566 6600  
2574 6620  
2582 6640  
2590 6660  
2598 6680  
2606 6700  
2614 6720  
2622 6740  
2630 6760  
2638 6780  
2646 6800  
2654 6820  
2662 6840  
2670 6860  
2678 6880  
2686 6900  
2694 6920  
2702 6940  
2710 6960  
2718 6980  
2726 7000  
2734 7020  
2742 7040  
2750 7060  
2758 7080  
2766 7100  
2774 7120  
2782 7140  
2790 7160  
2798 7180  
2806 7200  
2814 7220  
2822 7240  
2830 7260  
2838 7280  
2846 7300  
2854 7320  
2862 7340  
2870 7360  
2878 7380  
2886 7400  
2894 7420  
2902 7440  
2910 7460  
2918 7480  
2926 7500  
2934 7520  
2942 7540  
2950 7560  
2958 7580  
2966 7600  
2974 7620  
2982 7640  
2990 7660  
2998 7680  
3006 7700  
3014 7720  
3022 7740  
3030 7760  
3038 7780  
3046 7800  
3054 7820  
3062 7840  
3070 7860  
3078 7880  
3086 7900  
3094 7920  
3102 7940  
3110 7960  
3118 7980  
3126 8000  
3134 8020  
3142 8040  
3150 8060  
3158 8080  
3166 8100  
3174 8120  
3182 8140  
3190 8160  
3198 8180  
3206 8200  
3214 8220  
3222 8240  
3230 8260  
3238 8280  
3246 8300  
3254 8320  
3262 8340  
3270 8360  
3278 8380  
3286 8400  
3294 8420  
3302 8440  
3310 8460  
3318 8480  
3326 8500  
3334 8520  
3342 8540  
3350 8560  
3358 8580  
3366 8600  
3374 8620  
3382 8640  
3390 8660  
3398 8680  
3406 8700  
3414 8720  
3422 8740  
3430 8760  
3438 8780  
3446 8800  
3454 8820  
3462 8840  
3470 8860  
3478 8880  
3486 8900  
3494 8920  
3502 8940  
3510 8960  
3518 8980  
3526 9000  
3534 9020  
3542 9040  
3550 9060  
3558 9080  
3566 9100  
3574 9120  
3582 9140  
3590 9160  
3598 9180  
3606 9200  
3614 9220  
3622 9240  
3630 9260  
3638 9280  
3646 9300  
3654 9320  
3662 9340  
3670 9360  
3678 9380  
3686 9400  
3694 9420  
3702 9440  
3710 9460  
3718 9480  
3726 9500  
3734 9520  
3742 9540  
3750 9560  
3758 9580  
3766 9600  
3774 9620  
3782 9640  
3790 9660  
3798 9680  
3806 9700  
3814 9720  
3822 9740  
3830 9760  
3838 9780  
3846 9800  
3854 9820  
3862 9840  
3870 9860  
3878 9880  
3886 9900  
3894 9920  
3902 9940  
3910 9960  
3918 9980  
3926 10000  
3934 10020  
3942 10040  
3950 10060  
3958 10080  
3966 10100  
3974 10120  
3982 10140  
3990 10160  
3998 10180  
4006 10200  
4014 10220  
4022 10240  
4030 10260  
4038 10280  
4046 10300  
4054 10320  
4062 10340  
4070 10360  
4078 10380  
4086 10400  
4094 10420  
4102 10440  
4110 10460  
4118 10480  
4126 10500  
4134 10520  
4142 10540  
4150 10560  
4158 10580  
4166 10600  
4174 10620  
4182 10640  
4190 10660  
4198 10680  
4206 10700  
4214 10720  
4222 10740  
4230 10760  
4238 10780  
4246 10800  
4254 10820  
4262 10840  
4270 10860  
4278 10880  
4286 10900  
4294 10920  
4302 10940  
4310 10960  
4318 10980  
4326 11000  
4334 11020  
4342 11040  
4350 11060  
4358 11080  
4366 11100  
4374 11120  
4382 11140  
4390 11160  
4398 11180  
4406 11200  
4414 11220  
4422 11240  
4430 11260  
4438 11280  
4446 11300  
4454 11320  
4462 11340  
4470 11360  
4478 11380  
4486 11400  
4494 11420  
4502 11440  
4510 11460  
4518 11480  
4526 11500  
4534 11520  
4542 11540  
4550 11560  
4558 11580  
4566 11600  
4574 11620  
4582 11640  
4590 11660  
4598 11680  
4606 11700  
4614 11720  
4622 11740  
4630 11760  
4638 11780  
4646 11800  
4654 11820  
4662 11840  
4670 11860  
4678 11880  
4686 11900  
4694 11920  
4702 11940  
4710 11960  
4718 11980  
4726 12000  
4734 12020  
4742 12040  
4750 12060  
4758 12080  
4766 12100  
4774 12120  
4782 12140  
4790 12160  
4798 12180  
4806 12200  
4814 12220  
4822 12240  
4830 12260  
4838 12280  
4846 12300  
4854 12320  
4862 12340  
4870 12360  
4878 12380  
4886 12400  
4894 12420  
4902 12440  
4910 12460  
4918 12480  
4926 12500  
4934 12520  
4942 12540  
4950 12560  
4958 12580  
4966 12600  
4974 12620  
4982 12640  
4990 12660  
4998 12680  
5006 12700  
5014 12720  
5022 12740  
5030 12760  
5038 12780  
5046 12800  
5054 12820  
5062 12840  
5070 12860  
5078 12880  
5086 12900  
5094 12920  
5102 12940  
5110 12960  
5118 12980  
5126 13000  
5134 13020  
5142 13040  
5150 13060  
5158 13080  
5166 13100  
5174 13120  
5182 13140  
5190 13160  
5198 13180  
5206 13200  
5214 13220  
5222 13240  
5230 13260  
5238 13280  
5246 13300  
5254 13320  
5262 13340  
5270 13360  
5278 13380  
5286 13400  
5294 13420  
5302 13440  
5310 13460  
5318 13480  
5326 13500  
5334 13520  
5342 13540  
5350 13560  
5358 13580  
5366 13600  
5374 13620  
5382 13640  
5390 13660  
5398 13680  
5406 13700  
5414 13720  
5422 13740  
5430 13760  
5438 13780  
5446 13800  
5454 13820  
5462 13840  
5470 13860  
5478 13880  
5486 13900  
5494 13920  
5502 13940  
5510 13960  
5518 13980  
5526 14000  
5534 14020  
5542 14040  
5550 14060  
5558 14080  
5566 14100  
5574 14120  
5582 14140  
5590 14160  
5598 14180  
5606 14200  
5614 14220  
5622 14240  
5630 14260  
5638 14280  
5646 14300  
5654 14320  
5662 14340  
5670 14360  
5678 14380  
5686 14400  
5694 14420  
5702 14440  
5710 14460  
5718 14480  
5726 14500  
5734 14520  
5742 14540  
5750 14560  
5758 14580  
5766 14600  
5774 14620  
5782 14640  
5790 14660  
5798 14680  
5806 14700  
5814 14720  
5822 14740  
5830 14760  
5838 14780  
5846 14800  
5854 14820  
5862 14840  
5870 14860  
5878 14880  
5886 14900  
5894 14920  
5902 14940  
5910 14960  
5918 14980  
5926 15000  
5934 15020  
5942 15040  
5950 15060  
5958 15080  
5966 15100  
5974 15120  
5982 15140  
5990 15160  
5998 15180  
6006 15200  
6014 15220  
6022 15240  
6030 15260  
6038 15280  
6046 15300  
6054 15320  
6062 15340  
6070 15360  
6078 15380  
6086 15400  
6094 15420  
6102 15440  
6110 15460  
6118 15480  
6126 15500  
6134 15520  
6142 15540  
6150 15560  
6158 15580  
6166 15600  
6174 15620  
6182 15640  
6190 15660  
6198 15680  
6206 15700  
6214 15720  
6222 15740  
6230 15760  
6238 15780  
6246 15800  
6254 15820  
6262 15840  
6270 15860  
6278 15880  
6286 15900  
6294 15920  
6302 15940  
6310 15960  
6318 15980  
6326 16000  
6334 16020  
6342 16040  
6350 16060  
6358 16080  
6366 16100  
6374 16120  
6382 16140  
6390 16160  
6398 16180  
6406 16200  
6414 16220  
6422 16240  
6430 16260  
6438 16280  
6446 16300  
6454 16320  
6462 16340  
6470 16360  
6478 16380  
6486 16400  
6494 16420  
6502 16440  
6510 16460  
6518 16480  
6526 16500  
6534 16520  
6542 16540  
6550 16560  
6558 16580  
6566 16600  
6574 16620  
6582 16640  
6590 16660  
6598 16680  
6606 16700  
6614 16720  
6622 16740  
6630 16760  
6638 16780  
6646 16800  
6654 16820  
6662 16840  
6670 16860  
6678 16880  
6686 16900  
6694 16920  
6702 16940  
6710 16960  
6718 16980  
6726 17000  
6734 17020  
6742 17040  
6750 17060  
6758 17080  
6766 17100  
6774 17120  
6782 17140  
6790 17160  
6798 17180  
6806 17200  
6814 17220  
6822 17240  
6830 17260  
6838 17280  
6846 17300  
6854 17320  
6862 17340  
6870 17360  
6878 17380  
6886 17400  
6894 17420  
6902 17440  
6910 17460  
6918 17480  
6926 17500  
6934 17520  
6942 17540  
6950 17560  
6958 17580  
6966 17600  
6974 17620  
6982 17640  
6990 17660  
6998 17680  
7006 17700  
7014 17720  
7022 17740  
7030 17760  
7038 17780  
7046 17800  
7054 17820  
7062 17840  
7070 17860  
7078 17880  
7086 17900  
7094 17920  
7102 17940  
7110 17960  
7118 17980  
7126 18000  
7134 18020  
7142 18040  
7150 18060  
7158 18080  
7166 18100  
7174 18120  
7182 18140  
7190 18160  
7198 18180  
7206 18200  
7214 18220  
7222 18240  
7230 18260  
7238 18280  
7246 18300  
7254 18320  
7262 18340  
7270 18360  
7278 18380  
7286 18400  
7294 18420  
7302 18440  
7310 18460  
7318 18480  
7326 18500  
7334 18520  
7342 18540  
7350 18560  
7358 18580  
7366 18600  
7374 18620  
7382 18640  
7390 18660  
7398 18680  
7406 18700  
7414 18720  
7422 18740  
7430 18760  
7438 18780  
7446 18800  
7454 18820  
7462 18840  
7470 18860  
7478 18880  
7486 18900  
7494 18920  
7502 18940  
7510 18960  
7518 18980  
7526 19000  
7534 19020  
7542 19040  
7550 19060  
7558 19080  
7566 19100  
7574 19120  
7582 19140  
7590 19160  
7598 19180  
7606 19200  
7614 19220  
7622 19240  
7630 19260  
7638 19280  
7646 19300  
7654 19320  
7662 19340  
7670 19360  
7678 19380  
7686 19400  
7694 19420  
7702 19440  
7710 19460  
7718 19480  
7726 19500  
7734 19520  
7742 19540  
7750 19560  
7758 19580  
7766 19600  
7774 19620  
7782 19640  
7790 19660  
7798 19680  
7806 19700  
7814 19720  
7822 19740  
7830 19760  
7838 19780  
7846 19800  
7854 19820  
7862 19840  
7870 19860  
7878 19880  
7886 19900  
7894 19920  
7902 19940  
7910 19960  
7918 19980  
7926 20000  
7934 20020  
7942 20040  
7950 20060  
7958 20080  
7966 20100  
7974 20120  
7982 20140  
7990 20160  
7998 20180  
8006 20200  
8014 20220  
8022 20240  
8030 20260  
8038 20280  
8046 20300  
8054 20320  
8062 20340  
8070 20360  
8078 20380  
8086 20400  
8094 20420  
8102 20440  
8110 20460  
8118 20480  
8126 20500  
8134 20520  
8142 20540  
8150 20560  
8158 20580  
8166 20600  
8174 20620  
8182 20640  
8190 20660  
8198 20680  
8206 20700  
8214 20720  
8222 20740  
8230 20760  
8238 20780  
8246 20800  
8254 20820  
8262 20840  
8270 20860  
8278 20880  
8286 20900  
8294 20920  
8302 20940  
8310 20960  
8318 20980  
8326 21000  
8334 21020  
8342 21040  
8350 21060  
8358 21080  
8366 21100  
8374 21120  
8382 21140  
8390 21160  
8398 21180  
8406 21200  
8414 21220  
8422 21240  
8430 21260  
8438 21280  
8446 21300  
8454 21320  
8462 21340  
8470 21360  
8478 21380  
8486 21400  
8494 21420  
8502 21440  
8510 21460  
8518 21480  
8526 21500  
8534 21520  
8542 21540  
8550 21560  
8558 21580  
8566 21600  
8574 21620  
8582 21640  
8590 21660  
8



FIGURE 2

1 ATGGAGGGCT GCATGGGGGA GGAGTCGTTT CAGATGTGGG AGCTCAATCG  
51 GCGCCTGGAG GCCTACCTGG GCCGGGTCAA GCGCTGGAG GAGCAGAATG  
101 AGCTGCTCAG CGCCGGACTC GGGGGGCTCC GCGACAATC CGCGGACAEC  
151 TCCTGGCGGG CGCATGCCGA CGACGAGCTG GCGGCCCTGC GTGCGCTCGT  
201 TGACCAACGC TGGCGGGAGA AGCACGCGGC CGAGGTGGCG CGCGACAACC  
251 TGGCTGAAGA GCTGGAGGGC GTGGCAGGCC GATGCGAGCA GCTGCGGCTG  
301 GCCCGGGAGC GGACGACGGA GGAGGTAGCC CGCAACCGGC GCGCCGTCGA  
351 GGCAGAGAAA TGCGCCCGGG CCTGGCTGAG TAGCCAGGGG GCAGAGCTGG  
401 AGCGcGAGcT AGAGGCTcTA CGCGTgGCGC ACGAGGAGGA GCGCGTCGGT  
451 CTGAACGCGC AGgCtgCCTG TGCCCCCCGc cTGCCcgGc cgcCCcgGcc  
501 tccccgcgCCG GCCCCGGAGG tAGAGGAGcT GGcAAGGCGA CTGGGCGAGG  
551 CGTGGCGCGG GGCAGTGCGC GGCTACCAGG AGCGCGTGGC ACACATGGAG  
601 ACGTCGCTGG ACCAGAcCCG CGAGCGCCTG GCCCCGGCGG TGCAGGGTGC  
651 CCGCGAGGTC CGCCTGGAGC TACAGCAGCT CCAGGCTGAG CGCGGAGGCC  
701 TCCTGGAGCG CAGGGCAGCG TTGGAACAGA GGTGGAGGG CCGCTGgcag  
751 GaGcGGcTGC GGGcLACTGA AAAGTTCCAG CTGGCTGTGG AGGCCCTGGA  
801 GCAGGAGAAA CAGGGCCTAC AGAGCCAGAT CGCTCAGGTC CTGGAAGGTC  
851 GGCAGCAGCT GGCGCACCTC AAGATGTCCC TCAGCCTGGA GGTGGCCACG  
901 TACAGGACCC TCCTGGAGGC TGAGAACTCC CGGCTGCAA CACCTGGCGG  
951 TGGCTCCAAG ACTTCCCTCA GCTTTCAGGA CCCAAGCTG GAGCTGCAAT  
1001 TCCCTAGGAC CCCAGAGGGC CGGCGTCTTG GATCTTTGCT CCCAGTCCTG  
1051 AGCCCAACTT CCCTCCCCTC ACCCTTGCCT GCTACCCTTG AGACACCTGT  
1101 GCCAGCCTTT CTTAAGAACC AAGAATTCCT CCAGGCCCGT ACCCCTACCT  
1151 TGGCCAGCAC CCCCATCCCC CCCACACCTC AGGCACCCTC TCCTGCTGTA  
1201 GATGEAGAGA TCAGAGCCCA GGATGCTCCT CTCTCTCTGC TCCAGACACA  
1251 GGGTGGGAGG AAACAGGCTC CAGAGCCCCCT GCGGGCTGAA GCCAGGGTGG  
1301 CCATTCTGTC CAGCGTCCTG CCTGGACCAG AGGAGCCTGG GGGCCAGCGG  
1351 CAAGAGGCCA GTACAGGCCA GTCCCCAGAG GACCATGCCT CCTTGGCACC  
1401 ACCCCTCAGC CcTGACCACT CCAGTTTAGA GGCTAAGGAT GGAGAATCCG  
1451 GTGGGTCTAG AGTGTTCAGC ATATGCCGAG GGGAAGGTGA AGGGCAAATC

FIGURE 2 (Continued)

1501 TGGGGGTTGG TAGAGAAAGA AACAGCCATA GAGGGCAAAG TGGTAAGCAG  
1551 CTTGCAGCAG GAAATATGGG AAGAAGAGGA TCTAAACAGG AAGGAAATCC  
1601 AGGACTCCCA GGTTCCTTTG GAAAAAGAAA CCCTGAAGTC TCTGGGAGAG  
1651 GAGATTCAAG AGTCACTGAA GACTCTGGAA AACCAGAGCC ATGAGACACT  
1701 AGAAAGGGAG AATCAAGAAT GTCCGAGGTC TTTAGAAGAA GACTTAGAAA  
1751 CACTAAAAAG TCTAGAAAAG GAAAATAAAA GAGCTATTAA AGGATGTGGA  
1801 GGTAGTGAGA CCTCTAGAAA AAGAGGCTGT AGGCAACTTA AGCCTACAGG  
1851 AAAAGAGGAC ACACAGACAT TGCAATCCCT GCAAAGGAG AATCAAGAAC  
1901 TAATGAAATC TCTTGAAGGT AATCTAGAGA CATTTTATT TCCAGGAACG  
1951 GAAAATCAAG AATTAGTAAG TTCTCTGCAA GAGAACTTAG AGTCATTGAC  
2001 AGCTCTGGAA AAGGAGAATC AAGAGCCACT GAGATCTCCA GAAGTAGGGG  
2051 ATGAGGAGGC ACTGAGACCT CTGACAAAGG AGAATCAGGA ACCCCTGAGG  
2101 TCTCTTGAAG ATGAGAACAA AGAGGCCTTT AGATCTCTAG AAAAAGAGAA  
2151 CCAGGAGCCA CTGAAGACTC TAGAAGAAGA GGACCAGAGT ATTGTGAGAC  
2201 CTCTAGAAAC AGAGAATCAC AAATCACTGA GGTCTTTAGA AGAACAGGAC  
2251 CAAGAGACAT TGAGAACTCT TGAAAAAGAG ACTCAACAGC GACGGAGGTC  
2301 TCTAGGGGAA CAGGATCAGA TGACATTAG ACCCCCAGAA AAAGTGGATC  
2351 TAGAACCACT GAAGTCTCTT GACCAGGAGA TAGCTAGACC TCTTGAAAAT  
2401 GAGAATCAAG AGTTCTTAAA GTCAC TCAA GAAGAGAGCG TAGAGGCAGT  
2451 AAAATCTTTA GAAACAGAGA TCCTAGAATC ACTGAAGTCT GCGGGACAAG  
2501 AGAACCTGGA AACACTGAAA TCTCCAGAAA CTCAAGCACC ACTGTGGACT  
2551 CCAGAAGAAA TAAATAAATC AGGGGGCAAT GAATCCTCTA GAAAAGGAAA  
2601 TTCAAGAACC ACTGGAGTCT GTGGAAGTGA ACCAAGAGAC ATTCAGACTC  
2651 CTGGAAGAGG AGAATCAGGA ATCATTGAGA TCTCTGGGAG CATGGAACCT  
2701 GGAGAATTG AGATCTCCAG AGGAGTAGAC AAGGAAAGTC AAAGGAATCT  
2751 GGAAGAGGAA GAGAACCTGG GAAAGGGAGA GTACCAAGAG TCACTGAGGT  
2801 CTCTGGAGGA GGAGGGACAG GAGCTGCCGC AGTCTGCAGA TGTGCAGAGG  
2851 TGGGAAGATA CGGTGGAGAA GGACCAAGAA CTGGCTCAGG AAAGCCCTCC

FIGURE 2 (Continued)

2901 TGGGATGGCT GGAGTGGAAA ATAAg gatga Ggcagagctg AatctAaGGg  
2951 agcaGgatGG CTTCACTGGG AAGGAGGAGG TGGTAGAGCA GGGAGAGCTG  
3001 AATGCCACAG AGGAGGTCTG GTTCCCAGGC GAGGGGCACC CAGAGAACCC  
3051 TGAGCCCAAA GAGCAGAGAG GCCTGGTTGA GGGAGCCAGT gTGAAGGGAG  
3101 GGgctgagGg cctCcaggac CCTgaAgGgC AATCACAACA GGTGGGGACC  
3151 CCAGGcCTCC AGGCTCCCCA GGGgctgCca gaGgcgatag agCCcctGgt  
3201 GgAagatgaT GTGGCCcCAG GGGGTGACca AGCCTCCCCA GAGGTCATGT  
3251 TGGGGTCAGA GCCTGCCATG GGTGAGTCTG CTGCGGGagC TGAGCCAGgc  
3301 ctGGGgcaGG GGgtGGgaGG gCtGGGGgaC CcaggCcATC TgACCAGGGA  
3351 AGAGGTGATG GAACCACCCC TGGAAGAGGA GAGTTTGGAG GCAAAGAGGG  
3401 TTCagGGCTT GGAAGGGCCT AGAAAGGACC TAGAGGAGGC AGGTGGTCTG  
3451 GGGACAGAGT TCTCCGAGCT GCCTGGGAAG AGCAGAGACC CTTGGgAGCC  
3501 TCCCAGGGAG GGTAGGGAGG AGTCagAGGC TGAGGCCCCC AGGGGAGCAG  
3551 AGGAGGCGTT CCCTGCTGAG ACCCTGGGCC AACTGGAAG TGATGCCCCCT  
3601 TCACCTTGGC CTCTGGGgTC AGAGGAAGCT GAGGAGGATG TACCACCAGT  
3651 GCTGGTCTCC CCCAGCCCAa CGTACACCCC GATCCTGGAA GATGCCCCCTG  
3701 GGCTrCAGCC TCAGGCTGAA GGGAGTCAGG AGGCTAGCTG GGGGGTGCAG  
3751 GGGAGGGCTG AAGCTGGGAa AGTAGAGAGC GAGCAGGAGG AGTTGGGTTC  
3801 TGGGGAGATC CCGAGGGGCC TCCAGGAGGA AGGGGAGGAG AGCAGAGAAG  
3851 AGAGCGAGGA GGATGAGCTC GGGGAGACCC TTCCAGACTC CACTCCCCCTG  
3901 GGCITCTACC TCAGGTCCCC CACCTCCCCC AGGTGGACCC CACTGGAGAG  
3951 CAGAGGCCAC CCCCTCAAGG AGACTGGAAA GGAGGGCTGG GATCCTGCTG  
4001 TCCTGGCTTC CGAGGGCCTT GAGGArCCCT CAGAAAAGGA GGAGGGGGAG  
4051 GAGGGAGAAG AGGAGTGTGG CCGTGA CTCT GACCTGTCAG AAGAATTGA  
4101 GGACCTGGGG ACTGAGGCAC CTTTTCTTCC TGGGGTCCCT GGGGAGGTGG

FIGURE 2 (Continued)

4151 CAGAACCTCT GGGCCAGgtg cCcCaGCTGC TACTGGATCC TGCAGCCTGG  
4201 GATCGAGATG GGGAGTCTGA TGGGTTTGCA GATGAGGAag AAAGTGGGGA  
4251 GGAGGGAGAG GAGGATCAGG AGGAGGGGAG GGAGCCAGGG GCTGGGCGGT  
4301 GGGGGCCAGG GTCTTCTGTT GGCAGCCTCC AGGCCCTGAG TAGCTCCCAG  
4351 AGAGGGGAAT TCCTGGAGTC TGATTCTGTA AGTGTGAGCG TCCCCTGGGA  
4401 TGACAGCTTG AGGGGTGCAG TGGCTGGTGC CCCCAGACT GCCCTGGAAA  
4451 CGGAGTCCCA GGACAGTGCT GAGCCTTCTG GCTCAGAGGA AGAGTCTGAC  
4501 CCTGTTTCCT TGGAGAGGGA GGACAAAGTC CCTGGCCCTC TAGAGATCCC  
4551 CAGTGGGATG GAGGATGCAG GCCCAGGGGC AGACATCATT GGTGTTAATG  
4601 GCCAGGGTCC CAACTGGAG GGGAAAGTCAC AGCATGTAAA TGGGGGAGTA  
4651 ATGAACGGGC TGGAGCAGTC TGAGGAAAGT GGGGCAAGGA ATGCGCTAGT  
4701 CTCTGAGGGA GACCGAGGGA GCCCCTTTCA GGAGGAGGAG GGGAGTGCTC  
4751 TGAAGAGGTC TTCGGCAGGG GCTCCTGTTC ACCTGGGCCA GGGTCAGTTC  
4801 CTGAAGTTCA CTCAGAGGGA AGGAGATAGA GAGTCCTGGT CCTCAGGGGA  
4851 GGAC

FIGURE 3

1 MEGCMGEESF QMWELNRRLE AYLGRVKALE EQNELLSAGL GGLRRQSADT  
51 SWRAHADDEL AALRALVDOR WREKHAAEVA RDNLAEELEG VAGRCEQLRL  
101 ARERTTEEVA RNRRAVEAEK CARAWLSSQG AELEERELEAL RVAHEEERVG  
151 LNAQAACAPR LPAPPRPPAP APEVEELARR LGEAWRGAVR GYOERVAHME  
201 TSLDQTRERL ARAVQGAREV RLELOOLOAE RGGLLERRAA LEORLEGRWQ  
251 ERLRATEKFQ LAVEALEQEK QGLOSQIAOV LEGRQOLAH L KMSLSLEVAT  
301 YRTILLEAENS RLQTPGGGSK TSLSFQDPKL ELQFPRTPEG RRLGSLLPVL  
351 SPTSLPSPLP ATLETPVPAF LKNQEFLOAR TPTLASTPIP PTPQAPSPAV  
401 DAEIRAQDAP LSLLQTOGGR KOAPEPLRAE ARVAIPASVL PGPEEPGGQR  
451 QEASTGQSPE DHASLAPPLS PDHSSLEAKD GESGGSRVFS ICRGEGEGQI  
501 WGLVEKETAI EGKVSSLOQ EIWEEEDLNR KEIQDSQVPL EKETLKSIGE  
551 EIQESLKTLE NQSHETLERE NOECPRSLEE DLETLSLEK ENKRAIKGCC  
601 GSETSRKRC RQKPTGKED TOTLQSLQKE NOELMKSLEG NLETFLFPGT  
651 ENQELVSSLO ENLESLTALE KENQEP LRSP EVGDEEALRP LTKENQEPLR  
701 SLEDENKEAF RSLEKENQEP LKTL EEEDQS IVRPLETENH KSLRSLEE QD  
751 QETLRITLKE TOORRRSLGE QDOMTLRPPE KVDLEPLKSL DQEIARPLEN  
801 ENQEFLLKSLK EESVEAVKSL ETEILESLSKS AGQENLET LK SPETQAPLWT  
851 PEEINKSGGN ESSRKGN SRT TGVCGSEPRD IQTPGRGESG IIEISGSMEP  
901 GEFEISRGVD KESORNLEEE ENLGKGEYQE SLRSLEEEO ELQPSADVQR  
951 WEDTVEKDQE LAQESPPGMA GVENKDEAEL NLREQDGFTG KEEVVEQGEL  
1001 NATEEVWFPG EGHPE NPEPK EORGLVEGAS VKGGAEGLOD PEGOSQOVGT  
1051 PGLQAPQGLP EAIEPLVEDD VAPGGDOASP EVMIGSEPAM GESAAGAEPG

## FIGURE 3 (Continued)

1101 LGQGVGGGSD PGHLTREEVM EPPLEESLE AKRVOGLEGP RKDLEEAGGL  
1151 GTEFSELPGK SRDPWEPPRE GREESAEAP RGAEEAFP AE TLGHTGSDAP  
1201 SPWPLGSEEA EEDVPPVLVS PSPTYTPILE DAPGLQPOAE GSQEASWGVQ  
1251 GRAEAGKVES EQEELGSGEI PEGLOEEGEE SREESEED EL GETLPDSTPL  
1301 GFYLRSP TSP RWTPLSRGH PLKETGKEGW DPAVLASEGL EEPSEKEEGE  
1351 EGEEECGRDS DLSEEFEDLG TEAPFLPGVP GEVAEPLGQV POLLLDPAAW  
1401 DRDGESDGFA DEEESGEEGE EDQEEGREPG AGRWGPSSV GSLQALSSSQ  
1451 RGEFLES DSV SVSVPWDDSL RGAVAGAPKT ALETESQDSA EPSGSEESD  
1501 PVS LEREDKV PGP LEIPSGM EDAGPGADII GVNGQGNLE GKSQHVNGGV  
1551 MNGLEQSEES GARNALVSEG DRGSPFQEEE GSALKRSSAG APVHLGQGQF-  
1601 LKFTQREGDR ESWSSGED